

TAXONOMIC INDEX

(Note: The letter "T" represents a reference appearing in a table and "F" is one in a figure or illustration.)

A

Abutilon theophrasti Medic., 207
Acalymma bivittatum (Kirsh), 266T
Acalypha spp., 208T, 209T
Acanthospermum hispidum DC., 208T, 209F
 Acaridae (family), 523
 Acarina (order), 260T, 271, 292
Achaea finita (Guenee), 268T
 Aceosejidae (family), 523
 Acrididae (family), 261T
Acrostylus patruelis H.S., 261T
Adelphia (genus), 293
Adelphocoris sp. nr. *apicalis* Reuter, 263T
Adoretus cribratus Harris, 265T
Adoretus umbrosus F., 265T
Aegocera rectilinea Boisduval, 268T
Afrius figuratus Germar, 263T
 Agaristidae (family), 268T
Agathis rubricinctus Ashmead, 297T
Agathis sp., 297T
Aglossa caprealis (Hübner), 521
Agnoscelis sp., 264T
Agonum spp., 303
Agrotis gurgistanus Faldernan, 266T
Agrotis ipsilon (Hufnagel), 268T
Agrotis repleta Walker, 268T
Agrotis segetum (Schiffmueller), 269T
Agrotis sp., 293
Ahasverus advena (Waltl), 521, 546, 546F, 547
 Aizoaceae, 346
Albizia jubibrissin, 178
Alicododes dentipes (Oliver), 267T
 Aleocharidae (family), 264T
Allodermes morogorensis Harris, 262T
Alphitobius daperinus (Panzer), 521, 549, 550F
Alphitophagus bifasciatus (Say), 522
Alternaria arachidis Kulk, 337
Alternaria alternata (Fr.) Keissler, 337
Alternaria sp., 334, 491
 Alydidae (family), 263T
Amara spp., 303
Amaranthus retroflexus L., 207
Amaranthus spp., 208T, 209F
Ambinervosae (section), 23T, 26F; chromosome complements, 30; genomic evolution, 41
Ambrosia spp., 208T
Amitermes evuncifer Silvestri, 262T
Amorbia emigratella Busch, 268T
Amphiploides (series), 22T, 41

Amphorophora (*Hyperomyzus*) *lactucae* (L.) 264T, 282
Amsacta albistriga (Walker), 268T; biology, 273; control of, 278, 281
Amsacta lactinea (Cremet), 273
Amsacta linola (Fabricius), 268T
Amsacta moorei (Butler), 268T; biology, 273; control of, 278
Anaemerus fuscus Oliver, 267T
Anarsia ephippias (Meyrick), 268T
Ancistrotermes crucifer (Sjostedt), 262T
Ancistrotermes latinotus (Holmgren), 262T
Anisodactylus spp., 303
Anisolabis (= *Euborellia*) *annulipes* (Lucas), 262T
Anisopteromalus calandrae (Howard), 523
 Annuae (series), 22T, 41
 Anobiidae (family), 523
Anomala antiqua Gyllenhal, 265T
Anomala atrovirens, 265T
Anomala phebaja Oliver, 265T
Anomala sp., 265T
Anoplocnemis curvipes F., 263T
Anthia sexguttata (F.), 307
 Anthicidae (family), 522
Anthicus cervinus La Ferté-Sénectère, 522
Anthicus ephippium La Ferté-Sénectère 522
Anthicus vicinus La Ferté-Sénectère 522
 Anthocoridae (family), 522
Anthrenus verbasci (L.), 522
Anticarsia gemmatilis Hübner, 269T; biology, 272-273; control of, 278-280; damage, 276
Antilocaris pilosulus (Stol), 523
Apanteles sp., 297T, 522
Aphanus (= *Naphius*) *apicalis* (Dallas), 263T
Aphanus sordidus (F.), 263T
Aphasmatylenchus straturatus, 384
Aphelenchoides arachidis, 384
Aphelenchoides spp., 384
Aphelenchus spp., 384
 Aphididae (family), 264T-265T
Aphis craccivora Koch, 264T; biology, 281-284; control of, 284-285, 308; damage, 284; research needs, 759; as a vector, 110-111, 281-284, 340-341, 344-346
Aphis critica (= *spiraeola*), 282
Aphis glycines Matsumura, 265T
Aphis gossypii Glover, 265T; as a vector, 282, 340, 345

Aphis laburni Kaltenbach, 264T
Aphis leguminosae Theob., 264T
Aphis spiraeola, 282, 344
Aphitobius laevigatus (F.), 522
Apiomerus crassipes crassipes (F.), 303
Apophyllia murina Gerstaecker, 267T
Apophyllia nigricollis Allard, 267T
Arachis (section), 22T, 26; aneuploidy, 30-31; archaeological evidence, 11; biological species concept, 24-25, 28; breeding strategies, 44; cell DNA, 29; chemotaxonomy, 29-30; chromosomes, 30, 32-39; cytogenetics, 30-41; cytology, 30-41; disease resistance, 43, 111; dispersion of, 12-15; drought resistance, 43, 106; early history, 1-11; flavonoids, 29; floral characters, 56-62; gene pools, 42-43, 45, 95-96; genetic divergence, 40; genomic evolution, 41-42; geographic range, 25; germplasm resources, 43, 95-96, 98-99; growing season, 14, 54; interspecific hybridization, 25; response to irradiance, 422; morphological characteristics, 21, 24; morphological species concept, 24; nucleic acids, 29; origin, 25, 27F, 95, 754; pest resistance, 43-45, 70-71, 106, 291-292; photosynthesis, 421-423; plant characters, 51-56; ploidy manipulation, 44-45; polyploidy, 30, 38; pod and seed characters, 63-68; qualitative characters, 52-72; quantitative characters, 72-85; species of, 21-23; taxonomic subdivisions, 22-23; wild species, 28, 34, 39, 43-44, 95-96, 101, 104, 291, 421, 423, 447, 450
Arachis angustifolia (Chod. et Hassl.) Killip, 21
 A. *batizocoi* Krap. et Greg., 22, chromosomes, 31F, 33F, 38; cytological differentiation, 33; genomic divergence, 41; interspecific hybridization, 27-28; irregular meiosis, 36; origin of cultivated peanut, 95
 A. *x batizogaea* Krap. et Fern., 22T, 29
 A. *bentharii* Handro, 22, 23T; chromosome number, 31T; resistance to thrips, 112
 A. *burkartii* Handro, 22, 23T; genomic evolution, 41-42; pepper spot and leaf scorch, 334
 A. *cardenasii* Krap. et Greg., 22T, 24, 39; aneuploidy, 31; chromosomes, 31T, 33F, 35F, 36F, 38; in linkage mapping experiments, 72; origin of cultivated peanut, 95; resistance of, 44, 107-108, 330
 A. *chacoense* Krap. et Greg., 22T, 24; chromosomes, 31T, 38, interspecific hybridization, 28; effect of pests, 285, 288; resistance of, 44, 107-108, 330, 343
 A. *correntina* (Burk.) Krap. et Greg., chromosomes, 31T, 32, 36F, 38-39; interspecific

hybridization, 25, 28, 35; culture of reproductive tissue, 448-449
 A. *diagoi* Hoehne, 21, 22T, 24; chromosomes, 30, 31T; interspecific hybridization, 25, 28
 A. *durandensis* Krap. et Greg., 22T, 24; chromosomes, 31T, 35F, 38-39; interspecific hybridization, 28, 35-36; effect of rhizobia, 444; effect of thrips, 288
 A. *glabrata* Benth., 21, 23T; chromosomes, 30, 31T; interspecific hybridization, 25, 28; pepper spot and leaf scorch, 334; effect of thrips, 288
 A. *guaranitica* Chod. et Hassl., 21, 22
 A. *hagenbeckii* Harms, 21, 23-24; chromosomes, 31T; interspecific hybridization, 28; pepper spot and leaf scorch, 334
 A. *belodes* (Martius) Krap. et Rig., 22T, 24; chromosome number, 31T, 39; interspecific hybridization, 28
 A. *hypogaea* L.: inheritance of albinism, 60T; aneuploidy, 30; and apparent photosynthesis, 421; response to CO₂ concentration, 423; cell DNA, 29; chromosomes, 30, 31T, 34, 39; composition, 655; effect of diseases, 285, 288, 333-334, 339, 341-342, 347; dispersion of, 13-15; early history, 1-11; epistatic gene interaction, 99; floral characters, 57; genetic variability, 50-52; geographic origin, 12-13; germplasm accessibility, 42, 95-96; and integrated pest management, 250 et passim.; interspecific hybridization, 35-37; and nitrogen fixation, 444; oil content, 114; origin, 1, 11, 13, 95, 754; photosynthetic rates, 104; reproductive tissue culture, 448; resistance of, 107-108, 112, 388; lack of resistance, 112-113; as seed parent, 25, 28, 35; subspecies of, 50-52, 64, 101; taxonomic classification, 21-22, 45, 50-51, 101, 642, 754; tryptophan content, 43; incompatibility with wild species, 98, 101
 A. *hypogaea* ssp. *fastigiata* Waldron, 50-51, 101; and dormancy, 642; fatty acid composition, 115; heterotic response, 78; inheritance of leaf color, 58; and scab, 336
 A. *hypogaea* ssp. *fastigiata* var. *fastigiata*, 51, 101; and dormancy, 642; hybrid vigor, 99
 A. *hypogaea* ssp. *fastigiata* var. *vulgaris*, 14, 51, 101; and dormancy, 642; and hybrid vigor, 99
 A. *hypogaea* ssp. *hypogaea*, 50-51, 101; and dormancy, 642; fatty acid composition, 115; heterotic response, 78; inheritance of leaf color, 58
 A. *hypogaea* ssp. *hypogaea* var. *hirsuta*, 12-14, 51, 101
 A. *hypogaea* ssp. *hypogaea* var. *hypogaea*, 50, 101; hybrid vigor, 99

- A. hypogaea* Te3, 346
A. ipaensis Greg. et Greg., 22T; chromosome number, 31T; interspecific hybridization 28
A. lignosa (Chod. et Hassl.) Krap. et Greg., 23T, 24; chromosome number, 31T; genomic evolution, 41
A. lutescens Krap. et Rig., 22, 23T; chromosome number, 31T
A. macedoi Krap. et Greg., 23T; chromosome number, 31T; resistance of, 112
A. marginata Gard., 21, 23; chromosome number, 30-31; interspecific hybridization, 28
A. martii Handro, 22, 23T
A. monticola Krap. et Rig., 22; cell DNA, 29; chromosome number, 31T; lack of cross-compatibility, 37; heterosis, 78; as gene pool, 42; interspecific hybridization, 28; mitotic chromosomes, 32F; pepper spot and leaf scorch, 334; wild species meiosis, 34
A. nambyquarae Hoehne, 21-23, 30; somatic complements, 34
A. oteroi Krap. et Greg., 23T; chromosome number, 31T
A. paraguayensis Chod. et Hassl., 221, 223T, 32; chromosome number, 31T
A. pintoi Krap. et Greg., 23T; genomic evolution, 41
A. pusilla Benth., 21, 23T; chromosome complements, 30; chromosome number, 31T; genomic evolution, 41; resistance of, 343
A. rasteiro, 34
A. repens Handro, 22, 23T; chromosome number, 31T; genomic evolution, 41; resistance of, 112
A. rigonii Krap. et Greg., 22, 23T, 24, 36; chromosome number, 31T; genomic evolution, 41
A. sp. 9646, 32
A. sp. GKP 9841, 28
A. sp. 9901 GKP, 28
A. sp. 10017, 33
A. spegazzinii Greg. et Greg., 22T, 36; chromosome banding pattern, 38; chromosome number, 31T; interspecific hybridization, 28
A. stenoperma Greg. et Greg., 22T, 36; chromosome banding pattern, 38; chromosome number, 31T; interspecific hybridization, 28; resistance of, 44
A. tuberosa Benth., 21, 23T
A. villosa Benth., 21-22; aneuploidy, 30; chromosome banding pattern, 38; chromosome number, 31T; interspecific hybridization, 28, 32
A. villosa var. *correntina* Burkart, 22T. See also *A. correntina*
A. villosulicarpa Hodhne, 21-22, 23T, 45; chromosome number, 30, 31T; interspecific hybridization, 25, 28; tryptophan content, 43
Arachnida (class), 259, 260T
Araecerus fasciculatus (De Geer), 521, 543-544, 545F
Araneida (order), 523
Arctiidae (family), 268T; biology, 272-274
Ascochyta adzamehica, 332
Ascochyta arachidis, 332
Ascotis reciprocaria Walker, 268T
Ascotis (= *Boarmia*) *selenaria* Schiffermueller, 268T
Aspavia armigera F., 264T
Aspergillus amstelodami (Mangin) Thom and Church, 493, 495
Aspergillus candidus Link ex Fr., 493
Aspergillus chevalieri (Mangin) Thom and Church, 494-495
Aspergillus elegans, 712F
Aspergillus flavus Link ex Fries, 146, 290, 351F, 386-387, 466, causal organism, 377-378, 496-505, 511-513; control of, 110, 378, 505-511, 513; damage, 116, 377; disease cycle & epidemiology, 378, 486-496, 513; effect on grading, 739, 744; research needs, 755, 761
Aspergillus fumigatus Fres., 493, 495
Aspergillus glaucus, 492-495
Aspergillus niger Van Theighem, 488, 505; causal organism, 367, 377; control of, 367-368; disease cycle & epidemiology, 367; occurrence, 490, 492-496, 513; symptoms, 366
Aspergillus oryzae (Ahlb) Cohn, 496, 511, 712F, 721
Aspergillus parasiticus Speare, 110; causal organism, 377-378, 496-505, 513; control of, 507; occurrence, 486-496, 513; research needs, 761; in storage, 492, 496-497
Aspergillus pulverulentus (McAlpine) Thom., 367
Aspergillus repens deBary, 493, 495
Aspergillus ruber (Konig, Spieck., and Bremer) Thom and Church, 493, 495
Aspergillus spp., 488, 490, 492, 512
Aspergillus tamaris Kita, 493, 495-496
Aspergillus terreus Thom, 493, 513
Aspergillus versicolor (Vuill.) Tiraboschi, 511
Aspilota sp., 522
Astigmatidae (family), 260T
Astylus atramaculatus (Blanchard), 266T
Atherigona sp., 270T
Atractomorpha crenulata (F.), 268T
Atronesia sp., 264T
Atta capiguara Gonçalves, 270T
Attagenus elongatulus Casey, 522
Attagenus megatoma (L.), 521, 539, 540F

- Austracris guttulosa* Walker, 261T
Austroagallia torrida Evans, 282, 347
Austroasca alfalfae (Evans), 264T
Autographa (= *Phyometra*) *gamma* (L.), 269T
Avena sativa L., 644
Axonomorphae (section), 330

B

- Bacillus musae*, 376
Bacillus nicotianae, 376
Bacillus solanacearum, 376
Bacillus spp., 307, 364, 369
Bacillus thuringiensis Berliner, 279, 566
Baptisia indigo, 208T
Barathra brassicae (L.), 269T
Barombia humeralis Lab., 267T
Beauveria bassiana, 566
Beauveria tenella, 566
Belonolaimus (genera), 378, 379F, 383-384
Belonolaimus gracilis Steiner, 383-384
Belonolaimus longicaudatus Rau, 383-384, 388
Bemisia gossypiperda Mistra & Lamba, 264T
Bemisia inconspicua Quaint, 264T
Bemisia tabaci (Gennadius), 264T
Beta vulgaris, 345
Blapstinus pratensis LeConte, 521
Blattella sp., 262T
Blattidae (family), 262T
Boarmia selenaria Schiffermueller, 268T
Bombyliidae (family), 297T
Botryodiplodia theobromae Pat., 487, 490, 492
Botrytis cinerea (Persoon) ex Fries, 374
Brachiaria platyphylla (Griseb.) Nash., 208T
Brachystegia spp., 306
Brachytrypes membranaceus Drury, 261T
Bracon hebetor Say, 522
Bracon mellitor Say, 297T
Braconidae (family), 297T, 522
Bradysia sp., 523
Bromeliaceae, 342
Bufo marinus, 307
Bufo melanostictus, 307
Buphonella nigrovioleacea Jacoby var. *metallica*, 267T
Buprestidae (family), 266T

C

- Cadra cautella* (Walker), 294
Caesonia spp., 264T
Cajanus cajan, 345
Caliothrips braziliensis Morgan, 262T
Caliothrips fumipennis (Bagnall and Cameron), 263T; damage, 287
Caliothrips indicus Bagnall, 263T, control, 288; damage, 287
Caliothrips spp., 287
Caliothrips sudanesis (Bagnall and Cameron), 263T; damage, 287
Calocoris angustatus Lethierry, 263T
Caloglyphus (general), 289, 290
Caloglyphus micheali, 290
Caloglyphus spp., 506, 523
Calonectria crotalariae (Loos) Bell and Sobers, 357-359; causal organism, 357-358
Calonectria sp., 365
Calonectria theae, 358
Calonectria theae var. *crotalariae*, 358
Camponotus maculatus F., 270T
Canavalia ensiformis, 344, 346, 347
Cantharidae (family), 266T
Capsicum chinense, 12
Carcinops pumilio Erichson, 522
Carcinops quatuordecimstriata Stephens, 522
Carpophilus dimidiatus (F.), 521, 544, 545F
Carpophilus freemani Dobson, 522
Carpophilus hemipterus (L.), 522
Carpophilus humeralis (F.), 522
Carpophilus marginellus Motschulsky, 522
Carpophilus mutilatus Erichson, 522
Carpophilus pallipennis (Say), 522
Carpophilus pilosellus Motschulsky, 522
Cassia, 340
Cassia obtusifolia L., 208T, 209F
Cathartus quadricollis (Guérin-Meneville), 521
Catorama sp., 523
Caulophilus oryzae (Gyllenhal), 521
Caulorhizae (section), 23T, 26F, 37; genomic evolution, 41; resistance to mites, 292
Cecidomyiidae (family), 269T
Celatoria bosqi Blanch, 303
Celatoria diabroticae (Shimer), 303
Celatoria (= *Chaetophleps*) *setosa* (Coquillett), 303
Cenchrus spp., 208T
Centrosema pubescens, 346
Cephalonomia tarsalis (Ashmead), 521
Cerambycidae (family), 266T
Ceratobasidium, 363
Cercopidae (family), 264T
Cercospora arachidicola Hori, 201, 326-330, 334, 337, 350F; causal organism, 327-328; control of, 329-330, 356; disease cycle & epidemiology, 328-329; resistance to, 44, 71, 80, 106-108; symptoms, 327
Cercospora canescens Ellis & Martin, 327
Cercospora sp., 255
Cercosporidium personatum (Berk. & Curt.) Deighton, 201, 326-330, 334, 337, 350F; causal organism, 327-328; control of, 329-330, 356; disease cycle & epidemiology, 328-329; resistance to, 44, 71, 80, 106-108; symptoms, 327
Chaetomium spp., 489, 491, 496
Chaetophleps setosa (Coquillett), 303
Chalara elegans Nag Raj and Kend., 371
Chalara (genus), 371
Chalaropsis sp., 371

- Chalcididae (family), 297T
Chauliognathus sp., 266T
Cheilomenes vicina Mulsant, 288
Chelonus elasmopalpi McComb, 297T
Chelonus insularis (= *texanus*) (Cresson), 297T
Chelonus texanus (Cresson), 297T
Chenopodiaceae (family), 343, 346
Chenopodium album L., 208T, 209F, 341, 344
Chenopodium amaranticolor, 341, 343, 345, 347
Chenopodium quinoa, 341, 343-344, 346-347
Cheyletidae (family), 523
Cheyletus sp., 523
Chloropidae (family), 270T
Choanephora sp., 339
Chondracris rosea DeGeer, 261T
Chrotogonus hemipterus Schaum, 261T
Chrotogonus rotundatus Kirby, 261T
Chrotogonus senegalensis Krauss, 261T
Chrotogonus trachypterus (Blanchard), 261T
Chrysodeixis chalcites Esper, 269T
Chrysolagria nerei Borchmann, 266T
Chrysomelidae (family), 266T-267T
Chrysopa sp., 288
Chrysopidae (family), 288
Cicadellidae (family), 264T
Cicadulina arachidis China, 264T
Cicadulina similis China, 264T
Cicadulina sp., 264T
Cladosporium spp., 496
Claviceps paspali Stevens and Hall, 512
Coccidae (family), 265T
Coccinelidae (family), 266T, 284, 288
Colaspis (= *Maecolaspis*) *jansseni* (Bechyne), 267T
Colemania sphenarioides Boliver, 261T
Coleoptera (order), 265T-268T, 271, 292, 301-302; biology, 302-303; control of, 303-305; damage, 303; as a parasite, 284, 303
Coleopterus unicolor Say, 522
Colletotrichum capsici (Syd.) But. & Bis., 335
Colletotrichum dematium (Pers. ex Fr.) Grove, 335
Colletotrichum mangenoti, 335
Colletotrichum spp., 335-336, 337-338
Compositae, 342-343
Conipoda calcarata Saussure, 261T
Conoderus sp., 266T
Conotelus stenoides Murray, 522
Coptotermes formosanus Shiraki, 262T
Corcya cephalonica (Stainton), 521, 528, 528F, 529
Coreidae (family), 263T
Corticaria sp., 522
Corticium solani (Prill. and Del.) Bourd. and Galz, 363
Corticium vagum Berk. and Curt., 363
Coryna apicicornis Guérin-Meneville, 266T
Coryna hermanniae F., 266T
Coryna lanuginosa Gerstaecker, 266T
Cotalpa lanigera L., 265T
Cratopus punctum (F.), 267T
Crator cuniculus Burmeister, 266T
Creontiades pallidifer (Walker), 263T
Creontiades pallidus (Rambur), 263T
Cretonotus transiens Walker, 268T
Criconemiods (genera), 378, 383
Cristulariella pyramidalis Waterman and Marshall, 337
Croton spp., 208T, 209F
Cryptolestes ferrugineus (Stephens), 521
Cryptolestes pusillus (Schonherr), 521, 535, 537, 538F
Cryptolestes trucidus (Grouvelle), 521
Cryptophagidae (family), 523
Cryptoserphus abruptus (Say), 523
Crytozemia cognata Marshall, 267T
Cucujidae (family), 521
Cucumis anguria L., 209F
Cucurbitaceae, 343
Curculionidae (family), 267T-268T
Cyamopsis tetragonoloba, 347
Cyclocephala immaculata Oliver, 265T
Cydnidae (family), 264T
Cydnus sp., 264T
Cylindrocladium crotalariae (Loos) Bell and Sobers, 351F, 387; causal organism 357-358, control of, 358-359; disease cycle & epidemiology, 358; resistance to, 80, 109; symptoms, 357
Cynaues angustus (LeConte), 521
Cynodon dactylon (L.) Pers., 147
Cyperus esculentus L., 208T, 209F
Cyperus rotundus L., 208T
Cyperus spp., 208T
Cyrtomenus ciliatus (Palisot de Beavois), 264T
Cyrtomenus mirabilis (Perty), 264T
Cyrtorhinus caricoides Ghauri, 263T
Cyrtorhinus rectangularis Ghauri, 263T

D

- Dactyloctenium aegyptium* (L.) Richter, 208T, 209F
Darluc filum, 322
Dasychira georgiana Fawe., 269T
Datura stramonium L., 208T, 209F, 347
Decatoma affinis Oliver, 266T
Delphacidae (family), 264T
Dereodus recticollis Marshall, 267T
Dereodus vagabundus Faust, 267T
Dermaptera (order), 262T, 292
Dermestes ater De Geer, 521
Dermestes maculatus De Geer, 521
Dermeestidae (family), 522
Derochrus brevicollis Serville, 266T
Desmodium tortuosum (Sw.) DC., 208T, 209F
Diabrotica balteata LeConte, 267T, 302

E

- Echinochloa crus-galli* (L.) Beauv., 208T
Eiphoroma dentator (F.), 297T
Elasmopalpus angustellus Blanchard, 293
Elasmopalpus lignosellus Zeller, 268T, 270T, 293; biology, 293-295; control of, 111, 296-301, 308, 391; damage, 258, 295-296, 503, 506; pest status, 309
Elateridae (family), 266T
Eleusine indica (L.) Gaertn., 208T, 209F
Empididae (family), 270T
Empoasca alceda Ross & Cunningham, 264T
Empoasca dolichi Paoli, 264T
Empoasca fabae Harris, 112, 131, 264T; control of, 305
Empoasca facialis Jacobi, 264T
Empoasca flavescens (F.), 264T

- Empoasca solana* De Long, 264T
Empoasca spp., 264T; control of, 391; damage, 287; pest status, 309
Empoasca sudaica Bergman, 264T
Enneobrips flavens Moulton, 112, 263T; damage, 286-287
Entomophthora fresenii Nowakowski, 290
Entomophthora spp., 259, 290
Entomopoxvirus, 297
Eotetranychus, 260T
Ephestia caustella (Walker), 521, 526, 527F, 528
Epicauda cinerea Forester, 266T
Epicauda immaculata Say, 266T
Epicauda maculata (Say), 266T
Epicauda pennsylvanica (DeGeer), 266T
Epicauda sericans LeConte, 266T
Epicauda spp., 266T
Epicauda vittata (F.), 266T
Epilachna corrupta Mulsant, 266T
Epilachna similis var. *assimilis* Mulsant, 266T
Epilachna varivestis Mulsant, 266T
Epilachna vigintioctopunctata F., 266T
Epinotia opposita Heinrich, 268T
Erectoides (section), 22T-23T, 24, 26F, 27F; genomic evolution, 41; hybridization, 32-33, 36-37; interspecific gene flow, 42; meiotic behavior, 36; resistance of, 292, 330
Eremotermes nanus, 262T
Ergana bicolor Jacoby, 267T
Eruinia nicotiana, 376
Erysiphe communis (Wallr.) Fr., 338
Erysiphe pisi (PC) SH-Amans, 338
Erythronoea tripunctula (Melichar), 264T
Escherichia coli, 705
Estigmene acraea (Drury), 268T; control of, 278
Estigmene unipunctata Hampson, 268T
Euborellia annulipes (Lucas), 262T, 522
Euborellia stali Dohrn, 262T
Eucelatoria armigera (Coquillett), 279
Eufallia uncostata (Belonia), 522
Eulepida masbona Arrow, 265T; biology, 306; control, 307; damage, 307
Eulophidae (family), 288
Euphyotrips minozii Bagnall, 286
Eupodidae (family), 260T, 289
Euponera senaarensis (Mayr), 270T
Euproctis fasciata Walker, 269T
Euremia daira (Godart), 269T
Eurhizomatosa (series), 23T, 26F, 27F; genomic evolution, 42
Eurotium amstelodami Mangin, 493
Eurotium chevalieri Mangin, 495
Eurotium repens deB., 493
Eurotium rubrum König, Spieck., and Bremer, 493
Euspilotus sp., 522
Eutermes parvulus Sjostedt, 262T

Euxesta notata (Wiedemann), 523
Euxoa temera (Hubner), 269T
Extranervosae (section), 23T, 26F, 27F; genomic evolution, 41; interspecific gene flow, 42; resistance of, 292, 330

F

Fannia canicularis (L.), 522
Feltia sp., 293
Feltia subterranea (F.), 269T; biology, 273; control of, 278; damage, 276
Ferrisia virgata (Cockerell), 265T
 Flatidae (family), 264T
Flavobacterium aurantiacum, 511
 Formicidae (family), 270T
Frankliniella fusca (Hinds), 112, 131, 263T; biology, 286; control of, 288, 305; damage, 286-287; as a vector, 282, 342
Frankliniella occidentalis Pergrauve, 263T; as a vector, 282, 287, 342
Frankliniella schultzei Trybom, 263T; control of, 288; as a vector, 282, 287, 342
Frankliniella spp., 391
Frankliniella tritici (Fitch), 263T
 Fulgoridae (family), 264T
Furcifer rufiventris (L.W.) Johnson, 297
Fusarium equiseti, 365
Fusarium heterosporum, 365
Fusarium moniliforme, 365
Fusarium oxysporum, 365-366
Fusarium roseum, 365-366
Fusarium semitectum, 365
Fusarium solani (Mart.) Sacc., 365; causal organism, 365; control of, 361-362, 366; pod breakdown, 359, 360, 365; in stored seed, 493-494, 504-505
Fusarium spp., 360, 364-366, 382, 387, 513; and aeration, 502; causal organism, 365; control, 366; disease cycle & epidemiology, 365-366; dominance of, 488-489; occurrence, 372, 490-491; in stored seed, 492, 495-496; symptoms, 365
Fusarium tabacinum, 365
Fusarium tricinum, 365

G

Gecko gecko, 307
 Gelechiidae (family), 268T, 272-274
Geocoris sp., 288, 297
 Geometridae (family), 268T
Geron aridus Painter, 297T
Gibberella, 365
Gibbium psyllioides (Czenpinski), 522
Glomus mossae, 150
Glycine max L., 345-347, 357
Glycyphagus sp., 523
Gnathocerus cornutus (F.), 521
Gnathocerus maxillosus (F.), 521

Gonocephalem simplex (F.), 266T
Gonocephalum sp., 266T
Gossypium hirsutum L., 141
Graphognathus imitator (Bohemen), 267T
Graphognathus leucoloma (Boheman), 267T
Graphognathus peregrinus (Buchanan), 267T
Graphognathus sp., 267T
Graphognathus striatus (Boheman), 267T
 Gryllidae (family), 261T
Gryllotalpa africana Pal., 261T
Gryllotalpa eryllotalpa var. *casta*, 262T
Gryllotalpa gryllotalpa L., 262T
Gryllotalpa hexadactyla Perty 262T
 Gryllotalpidae (family), 261T-262T

H

Habrobracon gelechiae (Ashmead), 297T
Hallirbotius africana Jacoby, 267T
Halticus minutus Reuter, 263T
Halticus tibialis Reuter, 263T
Haplothrips gallarum Priesner, 263T; 287
Haplothysanus chapellei, 260T
Haplothysanus ealanus Attems, 260T
Haplothysanus haplothysarioides, 260T
Haplothysanus oubanguiensis Pierrard, 261T
Haptonicus luteolus (Erichson), 522
Hedylepta (= *Lamprosema*) *indicata* (F.), 268T
 Heleomyzidae (family), 523
Heliobis armigera (Hübner), 269T; biology, 272; control, 278
Heliobis depeacea (L.), 269T
Heliobis peltigera, 269T
Heliobis punctigera Walker, 269T
Heliobis spp., 273
Heliobis virescens (F.), 269T; biology, 272; control, 279
Heliobis zea (Boddie), 269T; biology, 272-273; control of, 278-280, 305, 391; damage, 276-278
Helicotylenchus dibystera (Cobb) Sher, 384
 Hemerobiidae (family), 288
Hemerobius sp., 288
 Hemiptera (order), 263T-264T, 271, 292
Hermetia illucens (L.), 522
Heterodera marioni, 380
Heteroligis claudius Klug, 265T
Heteronyx brevicollis Blackburn, 265T; biology, 306; damage, 307
Heteronyx spp., 306-307
Hippelates pusio Loew, 270T
Hippotion celerio (L.), 269T
Hister coenosus Erichson, 522
 Histeridae (family), 522
Hodotermes massambicus (Hagen), 262T
 Hodotermitidae (family), 262T
Holotrichia consanguinea (Blanchard), 265T
Homala sp., 266T
Homoneura sp., 270T
 Homoptera (order), 264T-265T, 271, 292

Howardula benigna Cobb, 303
Hylemya platura (Meigen), 523
 Hymenoptera (order), 270T, 271, 292
Hypomyces, 365
Hyperomyzus lactucae (L.), 264T, 340
Hypsonotus sp., 267T

I

Ichneumonidae (family), 297T, 522
Ilattia octo (Guenee), 269T
Indigofera endecaphylla, 512
Indigofera hirsuta, 208T
 Insecta (class), 259, 261T-270T
Invreia deceptor Grissell & Schauf, 297T
Invreia mirabilis (Boucek), 296
Invreia sp., 296
Invreia threa Grissell & Schauf, 297T
Invreia usta Grissell & Schauf, 297T
Ipomoea hederacea (L.) Jacq., 207, 209F
Ipomoea purpurea (L.) Roth., 207, 208T, 209F
Ischiodon aegypticus Wied, 288
Ischnotrachelus sp., 267T
Isodon puncticollis Maclure, 266T, 306
 Isoptera (order), 262T, 292

J-K

Jacquemontia tamnifolia (L.) Griseb., 208T, 209F
 Julida (order), 260T-261T
Kraussaria angulifera (Krauss), 261T

L

Labiduridae (family), 262T
Lachnosterna caudata; 265T
Lachnosterna (= *Holotrichia*) *consanguinea* (Blanchard), 265T; biology, 305-306; control of, 307; damage, 307
Lachnosterna fissa (Brenske), 265T
Lachnosterna spp., 307
Lactobacillus acidophilus, 722
Lactobacillus bulgaricus, 722
Lagria villosa F., 266T
 Lagriidae (family), 266T
Lasiobolus pallidulus Reuter, 522
Lasioderma sericorne (F.), 521, 537, 538F
Lateticus oryzae Waterhouse, 521, 535, 536F
 Lathridiidae (family), 522
 Lauxaniidae (family), 270T
 Lepidoptera (order), 268T-269T, 270, 292; control, 278-281, 308; as foliage consumers, 271-281; as soil-inhabiting pests, 293-301
 Leguminosae, 339, 346
 Leptidae (family), 270T
Leptosphaerulina arachidicola, 334, 336

Leptosphaerulina crassiasca (Sechet) Jackson & Bell, 334-337, 350F; causal organism, 334; control of, 335; disease cycle & epidemiology, 334-335; symptoms, 334
Leucopholis (genus), 307
 Limacodidae (family), 268T
Liogryllus morio F., 261T
 Liparidae (family), 269T
Liposcelis bastrychophilus Badonnel, 551
Liposcelis divinatorius (Muller), 551
Liposcelis spp., 521, 549, 551-552
Liurgus balteatus LeConte, 523
Locris sp., 264T
Locusta migratoria migratorioides Reiche and Fairmaire, 261T
Longidorus siddiqii, 387
Longunguis sacchari (Zehntner), 265T
Lophocarenum pusillum (Klug), 521
Loxostege strictalis L., 268T
Loxostege verticalis L., 268T
Luperodes quarternus Fairmaire, 267T
 Lycaenidae (family), 269T
 Lygaeidae (family), 263T, 288, 523
Lygaeus rivularis Germar, 263T
Lysiphebus fabarum (Marshall), 285
Lysiphebus testaceipes (Cresson), 284

M

Macrocentrus sp., 297T
Macrophoma sp., 334
Macrophomina spp., 368-369, 490
Macrophomina phaseoli (Maubl.) Ashby, 368, 487-489; causal organism 368-369; control of, 369; disease cycle & epidemiology, 369; occurrence, 490-491; symptoms, 368
Macrophomina phaseolina (Tassi) Goid., 487, 492, 513; causal organism, 368; disease cycle & epidemiology, 369
Macroposthonia ornata, 279F, 383-385, 387
Macroposthonia (genera), 378, 383, 385; control of, 391; population dynamics, 385
Macrosporium spp., 338
Macrotermes bellicosus (Smeathman), 262T
Macrotermes natalensis Haviland, 262T
Maecolaspis jansseni (Bechyne), 267T
Mamestra (= *Barathra*) *brassicae* (L.), 269T
Maruca testulalis (Geyer), 268T
Megachile argentata F., 270T
 Megachilidae (family), 270T
Megacoele stramineum Walker, 263T
Melanophthalma distinguenda (Comolli), 522
Melanoplus bivittatus (Say), 261T
Melanoplus differentialis (Thomas), 261T
Melanoplus femurrubrum (De Geer), 261T
Melanoplus sanguinipes (F.), 261T
Melichares sp., 523
 Meloidae (family), 266T

Melioidogyne arenaria (Neal) Chitwood, 379F, 385-386; control of, 113, 388-390, 392; interaction with mycotoxins, 290, 387; population dynamics, 385; symptoms, 380-381

Melioidogyne hapla Chitwood, 380-381, 384; control of, 389, 391; interaction with mycotoxins, 386-387; symptoms, 381

Melioidogyne incognita, 380

Melioidogyne javanica (Treub) Chitwood, 380, 388

Meloidogyne spp., 378, 380-381, 385; control of, 388; population dynamics, 385; sampling & detection, 386

Melyridae (family), 266T

Menida loriventris Germar, 264T

Mesoleurus dentipes (Marshall), 267T

Mesoplatys cincta Oliver, 267T

Mesostenus gracilis Cresson, 522

Messor spp., 270T

Metarrhizium anisopliae, 307, 566

Microcerotermes (= *Microtermes*) *parvulus* Sjöstedt, 262T

Microplitis croceipes (Cresson), 279, 297T

Microtermes parvulus Sjöstedt, 262T

Microtermes sp., 262T, 506

Microtermes thoracalis Sjöstedt, 262T

Miridae (family), 263T, 288

Mirperus jaculus Thunberg, 263T

Mocis repanda (F.), 269T

Mocis undata (F.), 269T

Molluga verticillata L., 208T

Monolepta australis (Jacoby), 267T

Monolepta goldinae Bryant, 267T

Monolepta nigeriae Bryant, 267T

Monolepta rosea Blackburn, 267T

Monolepta sp. nr. *kraatzii* Jacoby, 267T

Monophleboides arachidis Vayssiere, 265T

Monomorium bicolor Emery, 270T

Mononychus planki (McGregor), 260T

Monotoma sp., 523

Mucor hiemalis, 712

Musca domestica L., 522

Muscidae (family), 270T

Mycelia sterilia, 363

Mycetophagidae (family), 523

Mycoidiplosis sp., 269T

Mycosphaerella arachidicola, 322

Mycosphaerella arachidis Deighton, 327

Mycosphaerella argentinensis, 332

Mycosphaerella berkeleyi W. A. Jenkins, 327

Mylabris pustulata Thunberg, 266T

Mylabris sp., 266T

Mylabris trifasciata Thunberg, 266T

Myllocerus discolor Boheman, 267T

Myllocerus sp., 267T

Myllocerus viridanus F., 267T

Mymaridae (family), 288

Myrothecium roridum Tode ex Fr., 337

Myzus persicae Sulzer, 265T, 344; as a vector, 282, 340

N

Naphius apicalis (Dallas), 263T

Naphius zavattarii (Mancini), 263T

Nasutitermes sp., 262T

Naupactus cinereidorsum Hulst, 267T

Nectria, 365

Nematocerus acerbus Faust, 267T

Neoplectana sp., 303

Neurospora sitophila, 712F, 721

Nezara pallidiconsperata Stal., 264T

Nezara viridula (L.), 264T

Nicotiana benthamiana, 347-348

Nicotiana clelandii, 345-346, 348

Nicotiana glutinosa, 348

Nicotiana rustica, 348

Nicotiana tabacum, 348

Nitidulidae (family), 522

Noctuidae (family), 268T-269T, 272-274

Nomadacris septemfasciata Serville, 261T

Nosema heterosporum, 566

Nosema invadens, 566

Nosema oryzaephili, 566

Nosema plodiae, 566

Nosema whitei, 566

Notoxus calcaratus Horn, 522

Nythobia sp., 297T

O

Odontopygidae (family), 260T-261T

Odontotermes anceps (Sjöstedt), 262T

Odontotermes badius (Haviland), 262T, 503

Odontotermes latericius (Haviland), 262T, 503

Odontotermes nilensis Emerson, 262T

Odontotermes obesus (Rambur), 262T

Odontotermes sp., 262T

Odontotermes vulgaris (Haviland), 262T

Oidium arachidis Chorin, 338

Olethreutidae (family), 268T

Oligonychus pratensis (Banks), 260T

Ootheca bennigseni Weise, 267T

Ootheca mutabilis Sahlberg, 267T

Orgilus elasmopalpi Muesebeck, 296-297, 297T

Orgilus nitidus Muesebeck, 297T

Orgilus sp., 297T

Orygia mixta Snell, 269T

Orius (genus), 288

Orosius ablicinctus Distant, 264T

Orosius argentatus (Evans), 264T

Orthoptera (order), 261T-262T, 271, 292

Orthracis sp., 261T

Oryzaephilus mercator (Fauvel), 521, 533, 534F, 535

Oryzaephilus surinamensis (L.), 521, 533

Otitidae (family), 523

Oxya velox (F.), 261T

Oxyctonia versicolor (F.), 265T

P

Pachnaeus azurescens Gyllenhal, 267T

Pachnaeus costatus Perroud, 267T

Pachnaeus litus (Germar), 267T

Pachaeus psittacus Olivier, 267T

Pachylophus sp., 270T

Paecilomyces fumoso-roseus, 566

Palorus ratzeburgi (Wissmann), 521

Palorus subdepressus (Wollaston), 521

Pangaeus bilineatus (Say), 264T

Pangaeus congruus (Uhler), 264T

Panicum dichotomiflorum Michx., 208T, 209F

Panicum texanum Buckl., 208T, 209F

Pantomorus glaucus (Perty), 267T

Paraplonobia sp., 260T

Parasa vivida (Walker), 268T

Paratrachodorus (N) *christiei* (Allen) Siddiqi, 384

Paspalum dialatum, 512

Paspalum notatum Flugge, 147

Pedaliaceae, 346

Pellicularia filamentosa (Pat.) Rogers, 363

Pellicularia rolfsii (Curzi) West, 348-349

Penicillium camemberti Thom, 511

Penicillium citrinum Thom, 492-493

Penicillium cyclopium Westling, 511

Penicillium funiculosum Thom, 488-490, 492-494, 504-505, 513

Penicillium martensii Biourge, 495

Penicillium rubrum Stoll, 490, 492-492, 504-505

Penicillium spp., 382, 387; dominance, 488-489, 491; occurrence, 490; in stored seed, 492-493, 495-496

Pentatomidae (family), 263T-264T

Pentatoma major (Duges), 260T, 289

Pentodon idiota Herbst, 265T

Perennes (series), 22T, 41

Peridontopyge conani, 261T

Peridontopyge perplicata, 261T

Peridontopyge rebescens, 261T

Peridontopyge schoutedeni Attem, 261T

Peridontopyge sp., 261T

Peridontopyge spinosissima (Silvestri), 261T

Pestalotiopsis arachidis Satya, 338

Petunia hybrida, 348

Phaedonia areata F., 267T

Pharaxonotha kirschi Reitter, 523

Phaseolus aureus, 347

Phaseolus coccineus L., 40

Phaseolus mungo, 341, 347

Phaseolus vulgaris, 40, 343-345, 347; French Bean, 346; "Topcrop" 340

Phenacoccus hirsutus (Green), 265T

Philophaga viridicollis LeConte, 297

Phoma arachidicola Marasas, Pauer, and Boermea, 108, 350F; causal organism, 332-333; control of, 333-334; disease cycle & epidemiology, 333; symptoms, 332

Phomopsis sojae, 337

Phomopsis sp., 334

Phyllophaga ephialda Say, 265T

Phyllophaga micans Knock, 265T

Phylloticta arachidis-bypogaea, 338

Phylloticta spp., 338

Physalis floridana, 341

Physalospora rhodina (Berk. and Curt.) Cooke, 370

Phytomanas solanaceae, 376

Phytometra (= *Plusia*) *orichalcea* F., 269T

Phytometra gama L., 269T

Phytomonas solanaceae, 376

Pieridae (family), 269T

Piezodorus pallens Germar, 264T

Piezodorus rubrofasciatus F., 264T

Pisum sativum, 347

Planococcus (= *Pseudococcus*) *lilacinus* Cockerell, 265T

Platyedema ruficornis (Sturm), 522

Platynota nigrocyana Walsingham, 278

Platystomatidae (family), 270T

Pleospora crassiasca, 334

Plodia interpunctella (Hübner), 294, 521, 523-524, 525F

Plusia acuta Walker, 269T

Plusia chalcites (Esper), 269T

Plusia limbirena Guenee, 269T

Plusia orichalcea F., 269T

Plusia signata (F.), 269T

Podagrica sp. nr. *dilecta* Dalman, 267T

Podagrus (= *Crator*) *cuniculus* Burmeister, 266T

Polygonum pennsylvanicum L., 208T

Poophilus sp., 264T

Popillia japonica Newman, 266T

Portulaca oleracea L., 208T

Pratylenchus brachyurus Godfrey, 112-113, 379F, 381-384; control of, 388, 391-392; infection & spread, 382-383; interaction with mycotoxins, 386-387; symptoms, 381-382

Pratylenchus leiocephalus, 381

Pratylenchus spp., 378, 381-383, 392; population dynamics, 385

Prioclypeus bosqui Hulst, 267T

Pristomerus spinator (F.), 297T

Proctotrupidae (family), 523

Procumbensae (series), 23T, 41

Procrizomatosa (series), 23T, 26F, 41-42

Prostephanus truncatus (Horne), 521

Protostrophus ocularius Marshall, 268T

Protostrophus birtiventris Marshall, 267T

Psallus (= *Pseudatomoscelis*) *seriatus* (Reuter), 263T

Psallus sp., 288

Pseudatomoscelis seriatus (Reuter), 263T

Pseudococcidae (family), 265T
Pseudococcus brevipes (Cockerell), 265T
Pseudococcus calceolariae Maskell, 265T
Pseudococcus lilacinus Cockerell, 265T
Pseudococcus solani, 265T
Pseudococcus sp., 265T, 293
Pseudoleria pectinata (Loew), 523
Pseudomonas solanacearum (E. F. Sm.), 376-377; causal organism, 376; control of, 377; disease cycle & epidemiology, 376-377; symptoms, 376
Pseudomythyrus ancilla (Walker), 303
Pseudopachybrachius vinctus (Say), 523
Pseudoplusia includens Walker, 269T
Psilcephala acuta Adams, 297
Psocoptera (order), 549, 551
Psophocarpus tetragonolobus, 346
Pteromalidae (family), 523
Pterostichus spp., 303
Ptinus clavipes (Panzer), 522
Puccinia arachidis Spegazzini, 330-332, 350F; control of, 286; resistance to, 71, 108
Pyemotes sp., 523
Pyemotidae (family), 523
Pyralidae (family), 268T, 272-274, 294
Pyralis farinalis (L.), 521
Pyrgomorpha cognata Krauss, 261T
Pyrgomorpha vigneaudi Guérin-Meneville, 261T
Pyromorphidae (family), 268T
Pyrrhocoridae (family), 263T
Pythium myriotylum Drechsler, 109-110, 145, 290, 303, 351F; causal organism, 359-360; control of, 361-362; disease cycle & epidemiology, 360-361; pod breakdown, 359-360; symptoms, 360
Pythium sp., 345, 359-361; control of, 361-362; disease cycle & epidemiology, 360-361
Pythium ultimum, 390

R

R. oligosporus, 712
Radopholus similis (Cobb) Thorne, 384
Reduviidae (family), 303
Rhaconotus sp., 522
Rhizobium spp., 180, 380-381, 390; research needs, 756
Rhizoctonia bataticola (Taub.) Butler, 368
Rhizoctonia solani Kuhn, 145, 351F, 362-364, 489; causal organism, 337, 363; control of, 361-362, 364, 390, 392; disease cycle & epidemiology, 363-364; pod breakdown, 359-360, 362-363, 503; symptoms, 362, 382
Rhizoctonia spp., 145-146, 362, 488; control, 126, 217; dominance, 491; in stored seed, 492-493
Rhizomatosa (series), 23T, 29-30, 39; genomic evolution, 41-42; hybridization, 36-37; resistance of, 330

Rhizophagidae (family), 523
Rhizopus spp., 375, 513; and aeration, 502; causal organism, 375; occurrence, 490; in stored seed, 492
Rhizopus arrhizus, 375
Rhizopus nigricans Ehr., 487, 493
Rhizopus oryzae, 375
Rhizopus stolonifer (Ehr. ex Fr.), 375, 487; control of, 375; disease cycle & epidemiology, 375; in stored seed, 495-496
Rhopaea magnicoris Blackburn, 266T, 306-307
Rhopalosiphum maidis (Fitch), 265T
Rhopalosiphum padi (L.), 265T; as a vector, 282, 340
Rhyarochromus littoralis Distant, 263T
Rhizopertha dominica (F.), 521
Richardia scabra L., 208T, 209F
Rivellia sp., 270T
Romalea microptera Palisot de Beauvois, 261T

S

Sabra centurio Pascoe, 266T
Salmonella, 705
Sancassania sp. (= *Caloglyphus*), 260T
Scapteriscus abbreviatus Scudder, 262T
Scapteriscus aetetus Rehn & Hebard, 262T
Scapteriscus vicinus Scudder, 262T
Scaptocoris castaneus Perty, 264T
Scaptocoris terginus Schioedte, 264T
Scatopsidae (family), 523
Scarabaeidae (family), 265T-266T, 305
Scatopse fuscipes Meigen, 523
Scelionidae (family), 288
Scenopinus fenestralis (L.), 523
Scenopinus glabrifrons (Meigen), 523
Scenopinus nubilipes Say, 523
Septicus insularis Roelofs, 268T
Schistocera gregaria Forskal, 261T
Schizonycha africana Cast., 266T
Schizonycha sp., 266T
Sciariidae (family), 523
Schirtothrips dorsalis Hood, 263T; damage, 287; as a vector, 282, 287, 342
Sclerotinia arachidis, 354
Sclerotinia minor Jagger, 72, 329, 351F; causal organism, 354-355; control of, 356-357; disease cycle & epidemiology, 355-356; symptoms, 354
Sclerotinia miyabeana, 354
Sclerotinia sclerotiorum DeBary, 109; causal organism, 354-355; control of, 356-357; disease cycle & epidemiology, 355-356; symptoms, 354
Sclerotinia spp., 354-355
Sclerotinia trifoliorum, 354-355
Sclerotium bataticola Taub., 487, 492, 495; causal organism, 368; dominance, 489; interaction with mycotoxins, 505

Sclerotium rolfsii Sacc., 145, 351F; causal organism, 348-349; control of, 240, 329, 352-354, 390, 392; and cultural practices, 217-218; disease cycle & epidemiology, 349, 352; interaction with nematodes, 387; symptoms, 348
Sclerotium sp., 491, 493
Scolia aureipennis, 307
Scolothrips sexmaculatus (Pergande), 286
Scydmaenidae (family), 265T
Sebirus expansus Signoret, 264T
Selenothrips rebrocinctus Giard, 263T
Selepa docilis Butler, 269T
Senna tetraphylla, 10
Septanichus sp., 260T
Sericothrips occipitalis Hood, 263T, 287
Setaria faberi Herrm., 207, 208T
Setaria lutescens (Weigel) Hubb., 207, 208T
Setaria spp., 208T
Setaria viridis (L.) Beauv., 207, 208T
Sida spp., 208T, 209F
Silidius apicalis Waterhouse, 266T
Sitona crinitus (Hulse), 268T
Sitona lineatus (L.), 268T
Sitophilus oryzae (L.), 521
Sitophilus zeamais Motschulsky, 521
Sitotroga cerealella (Olivier), 521
Soja max, 346
Solanaceae (family), 342-343, 346
Solanum carolinense L., 208T, 758
Solanum elaeagnifolium Cav., 208T
Solenopsis fugax Latreille, 270T
Sorghum halepense (L.) Pers., 208T
Sorghum vulgare, 389
Spanagonicus albofasciatus (Reuter), 263T
Sphaceloma arachidis Bitancourt & Jenkins, 336
Speophtheria perroteti Guérin, 266T
Sphingidae (family), 269T
Sphrigodes globulus Marshall, 268T
Spilosoma investigatorium Karsch, 268T
Spodoptera eridania Cramer, 269T
Spodoptera exempta (Walker), 269T
Spodoptera exigua (Hübner), 269T, 272, 294; control of, 278-279; damage, 277-278
Spodoptera frugiperda Smith, 269T, 272, 294; control of, 278-281; damage, 274, 276-277; resistance to, 111-112
Spodoptera latifascia (Walker), 269T
Spodoptera littoralis (Boisduval), 269T
Spodoptera litura (F.), 269T, 272, 278
Spodoptera ornithogalli (Guenée), 269T, 272, 277-278
Spodoptera sp., 272
Sporomyxa tenebrionis, 566
Staphylinidae (family), 265T
Stegasta bosqueella (Chambers), 268T, 273-274; control of, 278, 391
Stegasta capitella (F.), 269T, 273-274

Stegobium paniceum (L.), 521, 537, 539, 540F
Stelidota chontalensis Sharp, 522
Stelidota geminata (Say), 522
Stelidota strigosa (Gyllenhal), 522
Stemphylium botryosum, 338
Stenopterochoris laticeps China, 263T
Stethorus, 291
Stomatomyia floridensis (Townsend), 297T
Stomatomyia parvipalpis (Wulp), 297T
Stomatomyia trinitatis (Thompson), 297T
Stomopteryx subsecivella (Zeller) (= *nerteria* (Meyrick)), 268T, 273; control of, 278, 280-281
Strigoderma arboricola F., 266T, 306
Strymon melinus (Hübner), 269T, 278
Stylopalpia costalimai Almeida, 268T, 293
Stylopalpia sp., 293
Stylosanthes biflora, 327
Stylosanthes spp., 283, 341
Syndesmogenus mimeuri, 261T
Syntermes sp., 262T
Syrphidae (family), 284, 288
Systates alticollis Marshall, 268T
Systates exaptus Marshall, 268T
Systates sp., 268T
Systema elongata F., 267T

T

Tachinidae (family), 297T
Taeniothrips distalis Karny, 263T
Taeniothrips inconsequens Uzel, 263T
Taeniothrips longistylus Karny, 263T
Taeniothrips sjostedi (Trybom), 263T; damage, 287
Tenebrio molitor L., 521, 547-549
Tenebrio obscurus F., 547
Tenebrionidae (family), 266T, 521
Tenebroides mauritanicus (L.), 541, 542F, 543
Termitidae (family), 262T
Tetrafoliatae (series), 23T
Tetramorium caespitum L., 270T
Tetramorium guineense (F.), 523
Tetraneura nigriabdominalis (Sasaki), 265T
Tetranychidae (family), 260T
Tetranychus arabicus Attiah, 260T
Tetranychus atlanticus McGregor, 260T
Tetranychus cinnabarinus (Boisduval), 258-259, 260T, 289; control of, 290-292
Tetranychus cucurbitacearum (Sayed), 260T
Tetranychus cucurbitae Rahman & Sapra, 260T
Tetranychus desertorum Banks, 260T
Tetranychus equatorius McGregor, 260T
Tetranychus hypogaeae, 260T
Tetranychus marianae McGregor, 260T
Tetranychus neocaledonicus Andre, 260T
Tetranychus spp., 260T, 291
Tetranychus telarius (L.), 260T

- Tetranychus tumidellus* Prichard & Baker, 260T, 292
Tetranychus turkestanii (Ugarov & Nikolski), 260T, 289
Tetranychus urticae Kock, 258-259, 260T, 289; control of, 290-292
 Tetrigidae (family), 261T
 Tettigometridae (family), 265T
Thanatephorus cucumeris (Frank) Donk, 363
Theobroma cacao, 345
Therioaphis ononidis (Kaltenbach), 265T
Thielaviopsis basicola (Berk. and Br.) Ferr., 371-372
 Thripidae (family), 262T-263T, 285
Thrips tabaci Lindeman, 263T, damage, 287; as a vector, 282, 342
 Thysanoptera (order), 262T-263T, 271, 285
Tibiomus ambitus (Attems), 261T
Tibiomus gossypii Pierrard, 261T
Tominotus communis (Uhler), 264T
Tominotus sp., 264T
 Tortricidae (family), 268T
Tortrix dinota Meyrick, 268T
Tota (genus), 293
Tribolium audax Halstead, 521
Tribolium castaneum (Herbst), 521, 530, 531F, 533
Tribolium confusum Jacquelin du Val, 521, 529, 531F
Trichoderma harzianum, 353, 364
Trichoderma lignorum, 353
Trichoderma spp., 353, 392; dominance, 489, 491
Trichoderma viride Pers. ex Fr., 353, 364, 369, 489, 496
Trichodorus christei Allen, 384
Trichodorus sp., 345
Trichogramma sp., 278-279
 Trichogrammatidae (family), 288
Trichonaupactus densius Hulst, 268T
Trichoplusia ni (Hübner), 269T, 278
Tricorynus ventralis (LeConte), 523
 Tridactylidae (family), 261T
 Trifoliolatae (series), 22T-23T
Trifolium repens, 344
Trigonidium cinctuloides Rambur, 261T
Trinervitermes biformis (Wasmann), 262T
Trinervitermes geminatus (Wasmann), 262T
 Triseminalae (section), 23T, 26F; chromosome complements, 30; genomic evolution, 41
Trissodon (= *Isodon*) *puncticollis* Maclure, 266T, 306-307
Triticum aestivum L., 41, 236
Trogium pulsatorium (L.), 551
Trogoderma glabrum (Herbst), 541
Trogoderma granarium Everts, 541
Trogoderma grassmani Beal, 541
Trogoderma inclusum LeConte, 521, 541, 542F
Trogoderma ornatum Say, 541
Trogoderma simplex Jayne, 541
Trogoderma sternale Jayne, 522, 541
Trogoderma sternale complex Casey, 522
Trogoderma sternale plagifer Casey, 522
Trogoderma teukton Beal, 541
Trogoderma variabile Ballion, 522, 541
Tuberculina costaricana, 332
Typhaea stercorea (L.), 521, 544, 546, 548
Tyrophagus sp., 260T, 290, 506, 523
- U**
- Ufa rebedinella* (Zeller), 293
Ufa (genus), 293
Uysanus sp., 264T
- V**
- Venturia canescens* (Gravenhost), 522
Verticillium albo-atrum Reinke and Berth., 372; casual organism, 373; control of, 373, 374; disease cycle & epidemiology, 373
Verticillium dahliae Kleb., causal organism, 373; control of, 373-374; disease cycle & epidemiology, 373; resistance to, 71
Verticillium spp., 110
Vigna sinensis Black Eye, 346
Vigna unguiculata, 341-343, 345-347
Voandzeia subterranea, 327
Vrydagba lepesmer, 265T
- W**
- Withius subruber* (Simon), 523
- X**
- Xanthium pensylvanicum* Wallr., 208T, 209F
Xanthium pungens Wallr., 213
Xantholinus spp., 303
Xanthomonas solanacearum, 376
Xiphinema diversicaudatum (Micoletzky) Thorne, 385
Xiphinema spp., 385
Xyletinus peltatus (Harris), 523
Xylocoris flavipes (Reuter), 522, 566
Xylocoris vicarius (Reuter), 522
Xylotrupes gideon L., 266T
Xyonysius californicus (Stol), 523
Xystrocerca marginalis F., 266T
- Z**
- Zea mays* L., 146, 236
Zinnia elegans, 348
Zoncerus elegans (Thunberg), 261T
Zoncerus variegatus F., 261T
Zophosis congesta Sjostedt, 266T
Zophosis sp., 266T
Zygrita diva Thomas, 266T

SUBJECT INDEX

A

- ABA. *See* abscisic acid
 Abnormal seedling, 418
 Aboveground field losses, 470
 Abscisic acid (ABA): and dormancy, 642-643; drought stress and stomatal closure, 178; growth regulation, 433-434; inhibition of germination, 416-417, 646; and pod formation, 437
 AC 3139, 107
 AC 91176, 114
 AC 91776, 114
 Acaricides, 291
 Acarines, 289
 Acetaldehyde, 636-637, 640
 Acetone, 636
 Acetone extraction, 656
 Acetyl value, 660T
 Acetylation, 712, 716
 Acids, 677T. *See also* names of individual acids
 Acifluorfen: nomenclature, 225T; relative toxicity of, 230T; technical description of, 224; weeds controlled by, 232T
 Acosta, Jose de, 4
 Acreage quota system, 737-738
 Acridines, 97
 Acriflazine, 70
 Activated charcoal, 238
 Actual evapotranspiration (E_a), 183; effect of climatic conditions, 189-191; effect of diseases, 191; effect of ground cover, 191.
See also Evapotranspiration
 Adaptation, 84
 Additional peanuts, 741, 744
 Additive epistatic variance, 99
 Additive genetic effects: breeding for early maturity, 114; inheritance of resistance, 80, 108; seedcoat splitting, 116; significance of, 80
 Additive genetic variance, 84; early generation testing, 85; importance of, 79, 99
 Adenosine phosphates, 80, 144
 Adenylate energy levels, 418
 Advanced breeding lines, 98
 Aeration, 502-503; after fumigation, 559-561
 Aeration method, 593
 Aeration systems, 580T, 581, 583-585
 Aerosol generator, 561-563
 Aerosol insect fogging systems, 745
 Aerosol space treatments, 554, 561-564
 Aflaroot, 377-378; control of, 378
 Aflatoxin B₁, 486
 Aflatoxin, 377-378, 486-513; as causal organism, 377-378; control of, 378, 505-508, 607, 707; damage, 377, 503-504, 745, 752; disease cycle and epidemiology, 378; and diseases in man, 486; grading and sorting procedures, 509-513, 596, 598-599, 618; growth periods, 582, 593, 607; and marketing agreements, 508, 590, 739; research needs, 761-762; resistance to, 110, 378, 506-507; symptoms of, 377
 "Aflatoxin-negative" certificate, 598
 Aflatoxin production, 496-503; effects of CO₂, O₂, and N₂, 502-503; relation to crop sequence, 504; effect of drought stress, 113, 466, 499; effects of microbial interaction, 504-505; effects of moisture and relative humidity, 497-500; preharvest and postharvest, 378, 466, 491, 499, 513; effects of temperature and time, 501-502
 Aflatrem, 512
 After-ripening, 643-644, 646
 Agar-gel electrophoresis, 696
 Age-injury relationship. *See* Temporal tolerance
 Aggregation, 720
 "Aging period," 640
 Agricultural Adjustment Act (1938), 737-738
 Agricultural Stabilization and Conservation Service (ASCS), 588, 740
 Agriculture and Food Act (1977), 738; (1978), 740
 Agroecosystem: functioning of, 250; as management unit, 251-252
 AH 1192, 199
 Air circulation patterns, 601
 Air classification liquid cyclone fractionation, 702, 705
 Air-dried seed/hull maturity index (DMI), 628-629
 Air flow, 582-583; of combine cleaning apparatus, 472; energy consumption, 482; pressure drops vs. fan requirements, 480; rate of, 475, 479-480; resistance of peanuts to, 479
 Air movement, 585
 Air temperature. *See* Temperature
 Air turbulence. *See* Wind
 Air velocity, 479, 480F
 Ajugose, 668

- AK 12-24, 199
 Alachlor: interactions with pesticides, 239-240; nomenclature, 225T; postemergence treatments, 225; preemergence treatments, 223; relative toxicity of, 230T; technical description, 224-225; weeds controlled by, 233T
 Alanap, 225T
 Alanine, 657T, 658T-659T, 701
 Alar-85 433
 Albina, 61
 Albinism, 59, 60T, 72
 Albumins, 655, 689-690, 692
 Alcohol dehydrogenase, 637
 Alcohols, 675T-677T
 Aldehydes, 636, 673, 675T, 677T
 Aldicarb, 391
 Aldrin, 303
 Alfalfa, 78
 Alimentary toxic aleukia (ATA), 486
 Aliphatic hydrocarbons, 676T-677T
 Alkanolamine salt, 226
 Alkylating agents, 97, 103
 Alkylpyrazine compounds, 673
 Almond moth, 527F; control of, 565-566; type of damage, 526, 528, description of, 526, life history, 526
 α, α, α -trifluoro-2,6-dinitro-*N, N*-di-propyl-*p*-toluidine, 222
 α -arachin, 655-656; amino acid composition, 700-701; extraction and fractionation, 690, 692; molecular weight, 698-699; polypeptide chains, 695-696; principle components, 695-701. *See also* Arachin, Protein
 α -conarachin: amino acid composition, 700-701, extraction and fractionation, 690, 692; molecular weight, 698-699; principle components, 695-701. *See also* Conarachin, Protein
 α -dicarbonyl compound, 672
 α -tocopherol, 682-683
 Alternaria leafspot, 337
 Alternative branching, 50
 Altitude, effect on growth, 441
 Aluminum (Al), 141, 150, 757
 Aluminum phosphide, 560
 Ambient air conditions, 481
 Ambient temperatures, 577-578, 585
 Amendoim (peanut synonym), 16
 Amendois (peanut synonym), 3-4
 American Peanut Exporting Company (APEC), 751
 AMI. *See* Arginine Maturity Index
 Amiben, 225T
 Amides, 217
 Amine salt, in formulation of 2,4-DB, 228
 Amino acetone, 672
 Amino acid composition, 658T-659T; inheritance of, 76; of peanut proteins, 656, 657T, 679; of protein fractions, 698-700
 Amino acids, 43, 115, 329, 507, 657-659; and curing, 636; and flavor, 671-672; mean composition, 659T
 Ammonia, 511, 601, 658T-659T, 761
 Ammonium polyphosphate, 156
 Ammonium salt, in formulation of chloramben, 226
 Ammonium sulfate, 639
 Amphidiploidy, 39
 Anaerobic condition, 637
 Analogs, peanuts used for: meat, 725; non-fermented cheese, 723-724
 Anchic (peanut synonym), 7, 9
 Aneuploidy, 30-31, 34, 755
 Angle iron bar cylinder, 609
 Angoumis grain moth, 521, 551
 Anhydrite (CaSO_4), soluble calcium source, 145
 Anhydrous ammonia, 511
 Anilines, substituted, 217
 Anion exchange resin, 238
 Annual growth habit, genetics of, 55. *See also* Growth habit
 Anthocyanin (red pigment), 57
 Anthracnose, 335; causal organism, 336; control of, 336; symptoms of, 335-336
 Antiauxins, 433
 Antifeeding compounds, 280
 Antinutritional factors, 763
 Aphids, 270, 281-282; biology, 282-284; type of damage, 284; management of, 284-285; worldwide pest status, 309T
 Aphid-transmitted viruses, 281-282; groundnut eyespot, 346; GRAV, 341; PGMV, 346
 Aphid vectors, 282T, 340-341, 347
 Apnea, 512
 Apollo (cultivar), 100
 Apomixis, sporadic, 28
 Apparent photosynthesis (AP): relation to CO_2 and O_2 concentrations, 423-424; effect of irradiance, 422; effect of leaf and plant age, 424-425; maximum ranges of, 420T, 421; effect of temperature, 421-422; effect of water stress, 422-423. *See also* Canopy AP, Photosynthesis
 Appressoria, 349
 Apricot kernel oil, 683
 Apteroous aphids, 341
 Aqueous alcohol leaching, 705
 Aqueous alkaline extraction, 706
 Aqueous ethanol extraction, 510
 Aqueous Isopropyl alcohol extraction, 510
 Aqueous system hydroperoxides, 639

- Aquifer levels, 761
 Aquiferous cells, 178
 Arabinose, 667-668
 "Arachide," 10
 Arachidic acid, 661T, 662T, 663T; mature vs. immature seed, 662; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T
 Arachin, 29, 655, 689 et passim.; composition of, 656; extraction and fractionation, 689-690; gel-forming properties, 719-720; purification, 692-693; seed protein, 29
 Arachin-P6, 696; amino acid composition, 701; molecular weights of, 697T
 Archaeological evidence, for cultivated peanut, 11
 Arctiids, 273
 Argentine (cultivar): amino acid composition, 658T; fatty acid composition, 662, 663T; genetic studies, 32F, 75-76, 86, 102; effect of herbicides, 241; milling quality, 592; effect of nitrogen, 148; resistance of, 109, 288, 304, 383, 388; effect of water deficit on seed, 173
 Arginine content, 656, 657T-659T, 700-701; inheritance of, 69; effect of maturation, 76, 657
 Arginine maturity index (AMI), 624, 626, 629-632, 667T
 Arm ratios, quotient of, 38
 Aroma, 673; origin, 636; of peanut flour bread, 726. *See also* Flavor
 Aromatic hydrocarbons, 677T
 Arrhenius equation, 475
 Arthropods: insect pests, 520, 523; as vectors, 282T; world list of, 260-270
 Artificial drying/curing, 466, 474-475; 743, 761; alternative systems, 483-484; bin or wagon drying, 478-479; energy consumption, 481-482; fan requirements, 480-481, heat requirements, 481; resistance of peanuts to air flow, 479; smoke drying, 3-4; specific heat, 477-478; effects of temperature and relative humidity, 475-477; thin-layer drying relationships, 475; and toxin-free seed, 501, 504
 Artificial hybridization techniques, 100-101
 Artificial hybrids, 96-97
 ASAE standard, 579
 Ascorbic acid, 638, 644
 Ash, 633, 655, 668, 705T; and cornmeal muffins, 728
 Ashy stem blight, 368
 Asiriya Mwitunde (cultivar), 291, 367, 506
 Asparagine, 369, 671
 Aspartic acid, 656, 657T-659T, 700-701; and flavor, 671
 Aspergillus blight, 366
 Aspergillus crown rot, 351F, 366; causal organisms, 367; control of, 367-368; disease cycle and epidemiology, 367; symptoms of, 366
 Aspertoxin, 512
 Aspiration, 553, 575T
 Aspiration machine, 572-573
 Aspirin, relative toxicity of, 230T
 Association of Official Analytical Chemists (AOAC), 418, 598
 Astigmatids, 289
 Atheromata, 683
 Atmospheric content, 648-649; and growth of fungi, 494, 496
 Atmospheric pressure, 648-649
 Atmospheric vapor pressure deficit (VPD), 176
 Atrazine, 241, 353
 Attractants, 251
 Auger conveying system, 472
 Aureus (chlorophyll-deficient mutant), 59-61
 Auto-catalysis, 646
 Automatic diffusion poropmeter, 176
 Autoxidation, 640
 Auxins (IAA), 433, 438; and dormancy, 642; and pegging, 435; in reproductive tissue culture, 449; effect on root growth, 446
 Available soil calcium, 139, 413. *See also* Calcium
 Available soil moisture, 137. *See also* Moisture, Soil moisture
 Available soil water, 192, 200. *See also* Irrigation, Soil water, Water
 Axial-flow fans, 480-481
- B**
- Ba. 186-28, 659T
 Backcrossing, 102-103; cultivar development and, 85; in heritability, 75; in ploidy level manipulation, 45
 "Backlegging," 581
 Bacteria, and insect control, 566
 Bacterial count, 705T
 Bacterial wilt, 326, 376; causal organism, 376; control of, 377; disease cycle and epidemiology, 376; symptoms of, 376
 Bagging material, 52, 599, 601, 648
 Bahiagrass, 147
 Bakery products, peanuts used for: breads, 726-728; cakes, 729; cookies, 728-729; doughnuts, sweet yeast products, 729
 Balan, 225T
 Bald seed, 115
 BAP. *See* Benzylaminopurine
 Barley, 113, 175, 359

- Barley yellow dwarf, 341
 Barnyardgrass, 208T
 Basagran, 225T
 Bauhini, Gaspard, 4, 5, 7-8
 Bauhino, 4, 9
 Baume sodium hydroxide (16₀), 510
 Beans, 506
 "Bed effect," 127
 "Beetknives," 219
 Beetles, 347, 521-523; individual descriptions of, 529-543, 544-547
 Beggarweed, Florida, 207, 208T, 211F; growth effects, 213; problem of control, 244-245; proclivity, 214, 215T, 216T, 218T; response to herbicides, 222-223, 225, 228, 232T
 Behenic acid, 661T, 662T, 663T, mature vs. immature seed, 662; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T
 Benefin, 222, 224, 235-236, 239; interactions with herbicides and pesticides, 239-240; nomenclature, 225T; relative toxicity of, 230T; and soil moisture, 223, 234; technical description, 225; weeds controlled by, 232T
 Benomyl, control of foliar diseases, 329, 331, 333, 353, 364; control of soilborne diseases, 364, 369, 372, 374; and spider mite outbreak, 291
 Bentazon, 224; nomenclature, 225T; relative toxicity of, 230T; technical description, 225-226; weeds controlled by, 232T
 Benzyladenine, 448
 Benzylaminopurine (BAP), 417, 645
 β -mercaptoethanol, 693
 β -sitosterol, 666
 Beverage and soup bases, peanuts used for, 723
 Beverage-type products, peanuts used for, 721-724
 Bills of lading, 749
 Bin drying, 478-479; and fungal invasion, 493
 Bins, 576, 577T; and insect infestation, 555
 Bioassay procedures, 386, 417
 Biochemical aging effect, 640
 Biochemical changes, and quality loss, 574
 Biological control, 308T, 310, 353, 759
 Biological nitrogen fixation. *See* Nitrogen fixation
 Biological species concept, 24-25
 Biology, of peanut pests, 520-521, 523 et passim.
 Biotin, 670T
 Birds: as predators, 307; in toxicity testing, 226, 229-230
 Bis(dimethylthiocarbamoyl) disulfide, 368
 Black carpet beetle, 540F; type of damage, 539; description, 539; life history, 539
 Black flour beetle, 521
 Black fungus beetle, 522
 "Black heart," 128, 129F
 Blackhull, 326, 371, 755; causal organism, 371; control of, 372; disease cycle and epidemiology, 372; symptoms of, 371
 Black larder beetle, 521
 Black nut, 368
 Black soldier fly, 522
 Blanching, 509, 640, 740, 745
 Blanco Rio II (cultivar), 336
 Blanco Santa Fe (cultivar), 336
 Blazer, 225T
 Bleckley (cultivar), 661
 B-nine, 433
 Boll weevil, 17
 Bonaparte, Lucien, 14
 Bonifay sand, 141
 Booklice, 551F; type of damage, 551-552; description, 551; life history, 551
 Boron: application rates, 129; critical levels, 155; deficiency of, 129, 153-154; as fertilizer, 154-155; importance of, 129, 760
 Boron content, 668T
 Botanical description, of peanut: early historical, 1-5, 7, 9-11; morphologically defined, 21
 Botrytis blight, 374; causal organism, 374; control of, 374; disease cycle and epidemiology, 374; symptoms of, 374
 "Botrytis shoot disease," 374
 Branching patterns, 50-55; alternative, 50, inflorescences, 50-51; inheritance of, 53; sequential, 50; effects of variation, 101
 Bran preparations, 353
 Bray-1 extractant: composition, 152; critical level, 152; soil response to, 152
 Bray-2 extractant: composition, 152; critical level, 152
 Brazilian peanut type, dispersion of, 13
 Breads, peanuts used for: cornbread, corn muffins, 728; specialty, 727-728; white and whole wheat, 726-727
 Breakfast cereals, peanuts used for, 730-731
 Breeders seed, 116-117
 Breeding objectives, 103-117, 144; to counteract costs of production, 103; to improve drought resistance, 98, 113; to enhance handling and processing characteristics, 95, 104; to speed maturity, 98, 113-114; to improve quality and appearance, 95, 104-105, 114-116; research needs, 754-755; to increase pest resistance, 104, 106-113, 508, 513, 761; to increase yields, 95, 103-104
 Breeding procedures: for cultivar development, 85, 95-99; development of, 77; and gene action, 77; significance of, 84-85
 Breeding strategies, for exploitation of wild species germplasm, 44

- Breeding systems, 99-100; backcrossing, 102-103; hybridization, 100-101; mass selection, 100-102; for mutations, 103; pure line selection, 100, 102-103; recombination, 100-101; research needs, 755; selection methods, 101
 "Bridging", 577-578
 Bristly starbur. *See* Starbur, bristly
 Broadhorned flour beetle, 521
 Broadleaf signalgrass. *See* Signalgrass, broadleaf
 Broadleaf weeds: control of, 135, 222-223, 226-227; effect on harvesting, 214; problem of, 244; and recirculating sprayer, 224
 Broadnosed grain beetle, 521
 Broad-sense heritability, 74-75
 Brokers, in marketing process, 748-749
 Bromelain, 710, 716-717, 720
 Brookston soil, 235
 Brown spider beetle, 522
 Brown spot, 350F
 Bucket elevator, 579, 581, 607
 Bud necrosis, 288-289, 342. *See also* Necrosis
 Buildings, storage. *See* Warehouses
 Bulk density, 579, 580T, 601
 Bulk drying process, 478. *See also* Curing, Drying
 Bulk holding tanks, 472, 552
 Bulk method, for cultivar development, 85
 Bulk population selection, for cultivar development, 101
 Bulk treatments, in storage, 554-555
 Bull nettle berries, 573T
 Bunch type cultivars: chromosomes of, 32; genotype x pesticide interactions, 113; growth habit, 52-53; effects of humidity and temperature, 441; resistance of, 367; susceptibility of, 369
 Bur gherkin, 210F
 Burlap bags: and insect infestation, 552; and sampling techniques, 599
 Buttermilk, peanuts used for, 722
 Butylate, 241
 Buyers, peanut: foreign, 742, 752; hedging techniques, 753; market risks, 751-753
 Bynum Runner (cultivar): amino acid composition, 659T; fatty acid composition, 661

C

C₃ crop species, 420
 C₃ photosynthetic pathway, 428
 C₃ photosynthetic type, 423
 C₄ plants, 424
 C501, 291
 C-55-437, 180-181
 C-59-127, 180
 Cacaguante, cachuate (peanut synonym), 9, 16
 Cadelle, 542F, type of damage, 543; description, 541; life history, 543
 Caffeic acid, 643
 Cakes, peanuts used for, 729
 Calcium, 141-146; 639; application methods, 142, 145; deficiency of, 128, 129F, 142-144, 172-174, 413; effect of disease incidence, 145; effect of drought, 172-173, 413; in pegging zone, 139, 144-145, 171; requirements of plant, 129, 144-145, 172-173, 760; effect on quality, 412-413; research needs, 756-757; soluble supplements, 145-146; sources of, 128-129, 139-141, 145-146; translocation in plant, 142; uptake of, 128, 172-173, 413; effect on yields, 128, 139-140
 Calcium concentration, 83, 86; combining ability for, 80; in foliage, 146; and imitation milk, 722
 Calcium content, 667T, 669T
 Calcium hydroxide, 511
 Calcium pantothenate, 447
 Calcium soil amendments, 145-146
 Calcium sulfate (CaSO₄), 172, 174
 Callus tissue, 447
 Campostrol, 666
 Cancer, 486
 Candy, peanuts used for, 17, 746
 Canopy apparent photosynthesis (AP): effects of CO₂ exchange, 426-427; irradiance, 426; leaf area index, 425-426; leafspot damage, 427; light interception, 427; seasonal changes, 427. *See also* Apparent photosynthesis, Photosynthesis
 Canopy cover: control of collar rot, 371; and crop evapotranspiration, 181, 183-185; density of, 349; photosynthesis and, 329, 425; research needs, 755; vs. soybean and cotton canopies, 207; and weed competition, 207; yield response to water use, 186-187
 Capital requirements, mechanization and, 458
 Capric acid, 660
 Caprylic acid, 660
 Captafol: suppression of aflaroot, 378; control of charcoal rot, 369; effect on Sclerotinia blight, 356; side effects, 329-330
 Captan, 368-369
 Carabids, 297
 Caramelization, 671
 Carbamate compounds, 291
 Carbaryl, 291
 Carbofuran, 391-392

- Carbohydrates, 257; composition, 655, 666-668; content of roots, 151; and flavor, 671; nutritional quality, 684; and physiological maturity index, 633; research needs, 755
- ¹⁴C distribution, and PMI, 633
- Carbon dioxide (CO₂): effect of concentration on AP, 423-424; in controlled atmosphere fumigation, 561; reduction of dormancy, 417; exchange, 426-427; fixation, 329; production, 413, 649-650
- Carbon dioxide uptake: and growth of *A. flavus*, 496, 502-503; and diffusive resistance, 174; and drought tolerant cultivar, 181; research needs, 762
- Carbon disulfide, 390
- Carbonyl compounds, 640-641, 647, 673, 674T
- Carboxin, 352-353, 364
- Carcinogenicity, 229, 390. *See also* Cancer
- Carding-type picker, 474
- Carlaviruses, 346
- Carmine spider mite, 258
- Carotene, 61, 627, 670-671
- Carotenoid pigments, 61, 671
- Carpetweed, 208T; response to herbicides, 232T; proclivity, 214, 215T, 218T
- Carrot motley dwarf, 341
- Casein, 679, 680T, 681F, 723
- Caseinate, 724
- Cast iron bar cylinder, 609
- Cast iron type sheller, 609, 610T
- Cast skin damage, 524, 528, 529, 533, 535, 537, 539, 541, 543, 544, 546, 547, 549
- Catechols, 670
- Cation exchange capacity, 242
- Cation exchange resin, 238
- Caulimoviruses, 348
- CCC. *See* Commodity Credit Corporation
- CCC, growth retardant, 433; effect on yields, 436
- Cell culture, 755. *See also* Tissue culture
- Cell DNA, 29
- Cellulose, 668
- Centrifugal fans, 480-481, 585
- Centrifugation, 706, 709
- CEPA, 417, 434, 645
- "Cephalin" fractions (liver lipid), 683
- Cercospora leafspot, 326-327; causal organisms, 327-328; control of, 329-330; disease cycle and epidemiology, 328-329; inheritance of resistance, 70; symptoms of, 327
- Cercosporidium leafspot, 326-327; causal organisms, 327-328; control of, 329-330; disease cycle and epidemiology, 328-329; symptoms of, 327
- Cercosporin, 329
- Cereals, peanuts used for, 730
- Chalimbana (cultivar), 100, 103, 111
- Chalky soils, 373
- Chapati, 727
- Charcoal rot, 368; causal organism, 368-369; control of, 369; disease cycle and epidemiology, 369; symptoms of, 368
- Chardakov densimetric dye technique, 179
- Cheese analogs, nonfermented, 723-724
- Chelated copper, 156
- Chelated iron, 156
- Chelated zinc, 158
- Chemical mutations, 97; to induce cytoplasmic mutations, 86; in dwarfism, 56; inheritability and, 70
- Chemical pesticide control, 758-759. *See also* Pesticides
- Chemicals: control of aflatoxins, 513; technical descriptions of herbicides, 224-228; control of nematodes, 126, 389-393; control of stem rot, 352-353; as suppressive agent, 252; weed control practices, 213; *See also* individual names of chemicals
- Chemical treatments, postharvest, 644
- Chico (cultivar), 71-72, 114, 133
- Chips, peanuts used for, 730
- Chloramben, 224, 236, 238; detoxification and soil pH, 238; interaction with nitrogen, 237; nomenclature, 225T; interaction with pesticides, 240; and seed quality, 241; early research, 222; soil moisture requirement, 226, 233-234; and soil organic matter content, 238; technical description, 226; toxicity of, 230T; volatilization, 236; weeds controlled by, 232T
- Chloramphenicol, 70
- Chlordane, 303-304
- Chlorina (chlorophyll-deficient mutant), 60
- Chlorine salts, 237
- Chlorogenic acid, 643
- Chlorophyll: concentration and water-stress, 178; inheritance of deficiencies, 59-61, 85; deficiency caused by low iron content, 156; effect of morphactins, 433
- Chlorophyll-deficient mutants: genotypes of, 61T; kinds of, 59-60; physiological relationships, 61-62. *See also* individual names of mutants
- Chlorophylls a and b, 61
- Chloropicrin, 389-390
- Chlorosis: due to aphids, 281; caused by foliar diseases, 338-339, 343, 345; caused by mineral deficiencies, 156-158; caused by soilborne diseases, 347-348, 354, 357, 365, 372-373, 376-377; due to teranychids, 289; due to thrips, 286
- Chlorothalonil: control of botrytis blight, 374; leafspot, 329-330; peanut rust, 331; sclerotinia blight, 356; web blotch, 333

- Chlorotic rosette, 340
- Chlorophoram, 220
- Choanephora leafspot, 339
- Cholesterol, 683
- Choline, 670T
- Choline chloride, 447
- Christensen's malt-salt medium, 494
- Chromatography, 417, 690, 693
- Chromium (alloy), 609
- Chromogene, 68
- Chromosomes: banding, 37-38; behavior, 34; doubling of complement, 98; meiotic, 38; morphology, 32-33, 37; number of, 30 et passim.; pairing of, 32-33, 39-40, 45. *See also* Genetics
- Chronological history, of peanuts, 1-18
- Chrysomelids, 301
- Cigarette beetle, 538F; type of damage, 527; description, 537; life history, 537
- Cis-N-[(1,1,2,2-tetrachloroethyl)thio]-4-cyclohexene-1,2-dicarboximide, 329
- Citrinin, 513
- Clay loam soil, 167, 193
- Clay percentage, in soil types, 238
- Clay soil, 183, 193, 226-227
- Cleaning, 572-574; in the combine, 472; research needs, 762. *See also* Shelling, Sorting
- Cleaning facility, 574F
- CLER flavor score method, 678
- Climatic conditions: effect on evapotranspiration, 189-191; effect on growth, 437-443; effect on herbicides, 231, 233-239; effect on nitrogen fixation, 150; and oil stability, 665
- Climatic requirements: optimum conditions, 125; rainfall, 125; temperature, 125
- Clover, 333, 344
- Clusius, 4-7, 9, 14
- Coastal bermudagrass, 147
- Cobalt, 155; effects on nitrogen fixation, 150, 155; and high pH soils, 150-151
- Cobb loamy sand, 194-195
- Cobo, Father Bernabé, 4-5, 8-9
- Coccinellids, 291
- Cocklebur, common, 208T, 210F; competitive ability of, 213; as foreign material, 573T; effect of herbicides, 223, 232T, 244; problem of, 243; proclivity, 214, 215T, 216T, 218T
- Coconut oil, 660
- Code of good practices, 601
- Coffee bean weevil, 545F; type of damage, 544; description, 543; life history, 543-544
- Co-isolates, 707
- Colchicine, 38-39, 98
- Cold storage, 647, 649. *See also* Refrigeration
- Cold tolerance, 415
- Coleoptera, 301-302; biology, 302-303; damage, 303; management of, 303-305
- Collar rot, 108-109, 366, 369-371
- Color: albinism, 59; other chlorophyll deficiencies, 59-61; of corolla, 57; of inner testa, 69; internal hull color, 626, 632; of leaf, 58-59; in linkage mapping experiments, 72; of raw peanut seed, 670-671; effect on sorting rate and frequency, 595; of testa, 58, 65-68, 632
- Color sorting, 575, 595, 618. *See also* Electronic color sorting, Handpicking, Sorting
- Colorado de Cordoba (cultivar), 102
- Column chromatographic technique, 673
- Combine. *See* Peanut combine
- Combine bulk bins, 473
- Combining: aboveground field losses, 470; alternative methods, 473-474; and *A. flavus* invasion, 498; energy consumption, 473; effect on germination, 413; and increased insect infestation, 552; mechanics of, 471-473; timing of operation, 470-471
- Combines: efficiency of, 135; injury to seed and weed control, 412; insect infestation, 553
- Comet (cultivar), 133, 188F; fatty acid composition, 662T; and lesser cornstalk borer, 111; oleate/linoleate ratios, 664T; resistance of, 300
- Commercial cultivars: attributes of, 105; breeding procedures, 85; coding of, 98; and control of nematodes, 388; research needs, 757. *See also* Cultivars
- Commercial peanut belt, 123, 124F
- Commercial production practices, 125-135, 412. *See also* Cultivation, Cultural Practices, Production
- Commercial-type sheller, 608F, 609T
- Commercialization of peanuts, 16-18
- Comminuted meats, 719
- Committee on Dietary Allowances, 682
- Commodity Credit Corporation (CCC): non-quota/additional peanuts, 743; oilstock, 750; provisions of, 740; purpose of, 738; surplus peanuts, 751
- Commodity treatments, 554; conventional fumigants, 559-561; malathion supplemental surface sprays, 557-558; modified atmosphere fumigants, 561; protectants, 555-556; surface sprays, 556-557; synergized pyrethrins supplemental surface sprays, 558-559
- Common cocklebur. *See* Cocklebur, common
- Common lambsquarter. *See* Lambsquarter, common
- Common purslane. *See* Purslane, common

- Communications technology, 764
- Competition: general principles of plant, 206-207; of weeds in peanuts, 207-208, 213-214
- Complementary gene action, 116
- Composition of peanuts: amino acids, 657-659; carbohydrates, 666-668; flavor, 671-678; minerals, 667-669; oil, 660-666; proteins, 655-657; research needs, 762-763; texture, 671; vitamins, 670; wholesomeness, 678-679
- Computer technology, 764
- Conagin *Macrocarpa* (cultivar), 658T
- Conarachin, 655 et passim., 689 et passim.; amino acids, 657; composition, 656; extraction and fractionation, 689-690; purification, 692-693
- Concealed damage, 129, 172, 588, 590. *See also* Damaged seed
- Concentrates, 511, 704-706
- Concrete silos, 576, 577T
- Condensation, 583-585, 601
- Confused flour beetle, 531F, 532F; type of damage, 529; description, 529; life history, 529
- Congo (cultivar), 55, 71
- Conservation, of natural enemies, 279
- Consumers: acceptance of peanut, 738-739; needs and desires, 95, 117; importance of texture to, 671
- Consumption of peanuts, 17; and price support system, 738-739
- Contact herbicides, 135. *See also* Herbicides
- Containers: for international shipping, 749; for packaging in U. S., 745
- Contamination, 520, 523 et passim.
- Continuous screw press, 702-703
- Continuous variation, 50
- Controlled atmosphere fumigation, 561
- Conventional fumigants, 559-561
- Conventional warehouse, 578T
- Conveyor belts, 599, 607
- Conveyor house warehouse, 584
- Cookies, peanuts used for, 728-729
- Copper, 155; deficiency of 155-156; in fungicides, 155
- Copper ammonium carbonate, 329-330
- Copper chelate, 156
- Copper content, 413, 668, 669T
- Copper hydroxide, 329
- Copperleaf, 208T, 211F; control of, 225, 228, 232T; proclivity, 214, 215T, 216T, 218T
- Copper sulfate, 155, 356-357
- Co-precipitated isolates, 707
- Cordoroy (cultivar), 421
- Coriante (cultivar), 628
- Corientes-3 (cultivar), 72
- Corn, 79, 236, 241, 245; and aflatoxin contamination, 486; 504, 506, 512-513; in crop rotation, 125, 146; 274, 359, 371, 389; as foreign material, 573T, 575T, 603, 618; and insect infestation, 543, 544, 546; weed competition in, 207; and zinc toxicity, 158
- Corn earworm, 391
- Corn oil, 660, 682-683
- Corn sap beetle, 545F; type of damage, 544; description, 544; life history, 544
- Corn syrup solids, 723
- Corolla color, 57
- Cost-benefit ratios: in disease control, 759; in irrigation systems, 757; in weed control, 758;
- Cost effectiveness, 761
- Cost, insurance, and freight (CIF), 749
- Cotton, 78, 141, 147, 167, 245; and aflatoxin contamination, 512; and boll weevil, 17; canopy comparisons, 207; in crop rotation, 125-126, 359, 371, 389; response to direct radiation, 189-190; effect of herbicides, 241; response to pesticides, 237, 239; second-order interactions, 83; response to water stress, 175; weed competition in, 206-207; and zinc toxicity, 158
- Cottonseed: extracts, 718; flour, 726; isolates, 720; oil, 660, 682-683; oil substrates, 639; products, 717-718
- Cottonseeds, 486, 513, 707
- Coultering, 460
- Coumarin, 643
- Countercurrent extraction equipment, 704
- Cowpea, 339-342; cross-inoculation group, 149
- Cowpea aphid, 344
- Cowpea meal, 724
- Cowpea mild mottle, 345; causal agent, 345-346; control of, 346; epidemiology, 346; symptoms and host range, 345
- Cowpea mild mottle virus (CMMV), 345-346
- Crabgrass, 208T; control of, 221, 232T, 244; and mechanical removal, 208; proclivity, 214, 215T, 218T; effect of time of removal, 213. *See also* Large crabgrass
- Cracking stage: application of alachlor, 224; dinoseb, 220, 226; diphenamid, 226; herbicide mixtures, 221-222, 225, 227; naptalam, 227; sesone or 2,4-DB with dinoseb, 221
- Cracking stage mixtures, 135
- Cradle hoist, 572
- Creeping indigo, 512
- Crescent factors, 58
- Critical moisture content (CMC), 649
- Crop evapotranspiration. *See* Evapotranspiration
- Crop management, 437. *See also* Cultural practices, Management strategies

- Crop maturity predictors: relative color evaluations, 626-628; indirect methods, 625-626; maturity methodologies, 624-635; physiological maturity methods, 632; research needs, 763; specific component quantitation, 630-632
- Crop-pest relationships, 253-257
- Crop residues: and aflatoxin contamination, 505-506; burial of, 126, 352; as foreign material, 575T; and irrigation systems, 201; and leafspot inoculum, 328-329; and lesser mealworm, 549; longevity of peanut rust, 331
- Crop rotation: and aflatoxin contamination, 505-506, 513; and disease control, 125-126, 329, 353, 357, 359, 362, 371, 377; and fertilizer utilization, 125; and preharvest insect control, 250-251, 301; and nematode control, 125-126, 388-389; and weed control, 126, 242-243
- Crop row spacing. *See* Row spacing
- Crop sequence, 490, 504
- Crop stress, 195
- Crop tillage. *See* Tillage
- Crop variety, 242-243. *See also* Variety selection
- Crop-water economy models, 189
- Crop water use: defined, 184; seasonal trends, 184-185; yield response, 186. *See also* Evapotranspiration, Water use efficiency
- Cross-compatibility, 36-37, 40, 42, 101
- Cross-pollination: artificial, 97; natural, 96-97; natural outcrossings, 96
- Croton, 208T, 210F; control of, 232T; proclivity, 214, 215T, 216T, 218T
- Crowfootgrass, 208T, 211F
- Crown rot, 351F, 366-368
- Crude fat percentage, 705T
- Crude fiber, 705T
- Crude protein content, 633, 655
- Crushing stock, 750-751
- Cryogenic-vacuum distillation, 641
- Cryoprecipitation, 691F
- Cucumber mosaic virus, 343
- "Cucumo viruses," 343
- Cultivar development: in breeding systems, 99; hybridization, 100-101; limiting factors, 99, 102; methods of, 85, 100-102; research needs, 755, 760; role of U. S. government, 738. *See also* Genetics, Germplasm
- Cultivars: rate of apparent photosynthesis, 421; coding of commercial, 98; interaction with environment, 658; genotype x pesticide interactions, 413; geographic dissemination, 51; maintenance of, 116-117; non-resistant, 366; 372, 374, 388; resistant, 327, 330, 332, 341, 378, 506-507; evaluation of stability, 83-84; effects of temperature, 441-442; traits of South American and African 96. *See also* names of individual cultivars
- Cultivated peanut: archaeological evidence, 11; breeding of, 95-116; evolution of, 40-41; origin of, 50-52, 95, 754; practical applications of genetic studies, 41; polyploidy, 30; protein and oil content, 43, 114-115
- Cultivation: land preparation, 126; response of peanut to mechanical, 207; seedbed preparation, 127, 133; damage to seed pod, 487; weed control, 135, 240
- Cultural control: of aphids, 285; of coleoptera, 305; of lepidoptera, 301; research needs, 759; spider mites, 292; thrips, 288-289; of weeds, 134-135, 207-208, 213, 242-244; of white grubs, 307
- Cultural practices: climate requirements, 125; crop rotation, 125-126; disease control, 341, 352, 364; effect on evapotranspiration, 192-193; fertilization, 128-129; geographic areas of, 123, 124F; growth regulators, 135-146; irrigation, 136-137; land preparation, 126-127; influence on physiology, 412; planting time and spacing, 133-134; and seed dormancy, 416; soil requirements, 123, 125; variety selection, 129-133; weed control, 134-135, 207-208, 213, 242-244. *See also* Cultivation, Production
- Cumulative pan evaporation (CPE), 195-196, 200
- "Cup," 62, 70, 103, 156
- Curculionids, 301
- Curing, 474, 624, 635; defined, 458; vs. "drying," 475; and enzymes, 636; equilibrium moisture content, 463-466; effect on fatty acid composition, 665-666; and fungal invasion, 486-487, 490-492, 513; effect on germination, 413; and lipids, 636; losses in the field, 470; moisture content definitions, 461-463; and respiration, 635-636. *See also* Drying
- Curing techniques: artificial curing, 3-4; and breeders seed, 116; in the field, 466-470; and milling process, 593; optimization of, 507-508; research needs, 761, 763; smoke drying, 3-4. *See also* Artificial drying, Stackpole, Windrow
- Cyanidin, 68
- Cyanocobalamine, 447, 670T
- Cyclodiene insecticides, 303
- Cyclopiazonic acid, 511-512
- Cylinders, 609; design of, 610T-612T, 613
- Cylinder-type picker, 474

- Cylindrocladium* black rot (CBR), 106, 109, 350F, 351F, 755, 759; causal organism, 357-358; control of, 358-359; and crop rotation, 126; disease cycle and epidemiology, 358; resistance to, 77, 80; symptoms of, 357
- Cysteine, 720
- Cytex, 436
- Cytogenetics, 30-41; practical applications of, 41-45; mutation research, 69-70; research needs, 755
- Cytokinins: relation to dormancy, 417, 642-643; growth promotion, 433, 436, 645-646
- Cytological differentiation, 33-34
- Cytological studies, 37
- Cytology, 30-41; practical application of, 41-45
- Cytoplasmic influence, on growth habit, 52-53
- Cytoplasmic inheritance, of plant characters, 85-87
- Cytoplasmic mutants, 86
- Cytoplasmic polyhedrosis virus, 566
- Cytotaxonomic classifications, 754

D

- Dagger nematodes, 385
- Daily water requirements, 137. *See also* Irrigation, Water
- Daily water use, 185. *See also* Irrigation, Water use efficiency
- Dairy products, peanuts used for, 721-724
- Dalliagrass poisoning, 512
- Damaged seed: and curing treatments, 475; and fungal invasion, 493, 503-504; insect control and resistance, 565; removal of, for seed stock, 603; standards established for, 597T, 599T; susceptibility to insects, 520; and windrow drying, 471. *See also* Concealed damage
- Damping-off: causal organisms, 359-360, 362-363, 365; resistance to, 361
- Darcy law, 184
- Dark plumules, 143, 174, 412-413
- Dark respiration, 422, 427
- Darou IV (cultivar), 110
- Dazomet, 390
- DBCP: control of nematodes, 389-391; control of *R. solani*, 364; negative effects, 353, 390; effect on yield, 240
- DCNA, 356, 374
- DD, 387, 389
- DDT, 303-304
- DEAE-cellulose chromatographic patterns, 690-693
- DEAE-Sephadex A-50, 639
- "Death watch," 551
- Decanal, 673
- Decay: in grading procedure, 590; of seed, 375
- Decortication, 503. *See* Shelling
- Deep plowing treatments, 352
- Defatted peanut flakes, 702, 725
- Defatted peanut flour, 702-704, 715; carbohydrates, 667; emulsion capacity, 715-717; extrusion processing, 724; foam formation, 718-719; flow diagram, 703F; nitrogen solubility, 709; quality specification, 705T; sugar contents, 688; use in peanut products, 721-731
- Defatted peanut grits, 702, 725
- Defatted peanut meal, 510-511, 681F; globulin extraction, 689; and lesser mealworm, 549; nutritional quality, 680; production of, 702-704; amount of protein, 693; trypsin inhibitors, 680
- Defoliation: causal organisms, 272, 274-275, 289, 329, 332, 334, 338, 362; and cessation of growth, 432; quantification of effects of, 255; sensitivity to, 256; and temporal tolerance, 255
- Dehydration, 497-498, 500
- de Laet, Jan, 4-7, 9
- de la Vega, Garcilaso, 1, 16
- de Léry, Jean, 3, 5
- Delivery process, 743-744
- Delta-T devices, 176
- Deodorized kerosene, 563
- Depressed flour beetle, 521
- Dermal LD₅₀, 229T
- DES. *See* Diethyl sulfate
- de Sahagún, Friar Bernardino, 4
- Desserts, frozen, peanuts used for, 723
- Desorption processes, 463, 466T
- Desorption isotherms, 464F, 465T
- Detached-leaf technique, 330
- Detection procedures: for aflatoxin contamination, 508-509; of CBR, 357-358; of insects, 762; of sclerotinia blight, 355
- Deterioration of seed, 495; control of field disease and, 412; role of field fungi, 492; from initial injury, 411; from insect damage, 520, 530; effect of storage conditions, 414
- Detoxification, 237-238, 761
- Development, of new cultivars, 17, 101-102, 116. *See also* Cultivar development
- Dew, 190
- Dew-point temperature, 475
- Dextrose, 369
- DHT, 200; and rust resistance, 108
- Diazinon, 304
- Dibromochloropropane, 240, 364
- Dicarbonyls, 640
- Dichloropropenes, 389
- Dichlorvos, 554, 562; residue tolerances, 562; in space treatments, 563
- Dieneamine, 672

- Diethyl(1,2-phenylene bis iminocarbonothioyl) bis (carbamate), 372
- Diethyl sulfate (DES), 70, 103
- Diethylenetriamine pentacetate-triethanolamine, 156
- Differential scanning calorimeter, 478
- Diffusion equation, 475
- Diffusion resistance (DR): in field environments, 176-178; and photosynthesis, 174-176
- Dig—wait-to-dig evaluation, 625-626
- Digger-shaker efficiency, 135
- Digger-shaker-inverter, 214, 460, 507
- Digging: losses during, 460; mechanics of, 460-461; and mycoflora, 486-491, 498, 513; predigging operations, 460; timing of, 458-460
- Diglycerides, and curing, 636
- Dihydro-O-methylsterigmatocystin, 512
- Dimethyl ether, 510
- Dimethylethylpyrazine, 672
- Dinitramine, 353, 359
- Dinitrophenol, 357
- Dinoseb, 219-224, 239; effectiveness, 240-241, 244, 353, 357, 359; effect of environment, 236-237; interactions with herbicides, 225, 227, 239-240; nomenclature, 225T; relative toxicity of, 230T; technical description, 226; weeds controlled by, 232T
- Dephenamid, 221, 224; interactions with herbicides, 240; nomenclature, 225T; relative toxicity of, 230T; technical description, 226-227; weeds controlled by, 232T
- Diplodia blight, 370
- Diplodia collar rot, 369; causal organism, 370; control of, 371; disease cycle and epidemiology, 370-371
- Diplodia dry rot, 370
- Diplodia wilt, 370
- "Dipping" 581
- Direct-beam radiation, 175, 177-178
- Direct fired gas heaters, 481
- Direct gas chromatographic method, 641
- Direct vapor exchange techniques, 184
- Directed management, 252. *See also* Crop management, Integrated pest management
- Dirt: as foreign material, 573T; methods of removal, 575T
- "Dirting" cultivation, 217, 240; control of stem rot, 352
- Dirt screen, 572
- Disaccharides, 415
- Disc electrophoresis, 29
- Disc-gel electrophoresis, 707
- Discontinuous variation, 50
- Disease: avoidance of, 331-332; effect of crop rotation, 125-126; and digging losses, 460; effect on evapotranspiration, 191;

- foliar, 326-348; and harvesting, 625-626 and irrigation scheduling 187-201; caused by nematodes, 378-385; quantification of effects of, 255; research needs, 759-760; resistance to, 106-111, 507; soilborne, 348-378; and storage effects, 412; relation to weed control, 217-219, 242
- Disease management: anthracnose, 336; aspergillus crown rot, 371; bacterial wilt, 377; blackhull, 372; botrytis blight, 374; charcoal rot, 369; cylindrocladium black rot, 358-359; diplodia collar rot, 371; fusarium diseases, 366; groundnut rosette, 341; early and late leafspot, 329-330; nematodes, 387-393; peanut clump, 345; peanut mottle, 340; peanut stunt, 344; pepper spot and leaf scorch, 335; pythium diseases, 361-362; rhizoctonia diseases, 364; rhizopus diseases, 375; rust, 331-332; sclerotinia blight, 356-357; stem rot, 352-354; tomato spotted wilt, 342-343; verticillium wilt, 373-374; web blotch, 333-334; yellow mold, aflaroot and aflatoxin, 378, 505-508, 513
- Disk harrows, 228
- Disulfoton, 240
- Dithiothreitol (DTT), 693, 698
- Diuron, 220
- Divalent salts, 708-709
- Diversity, plant character associations, 73-75
- Dixie Anak (cultivar), 71
- Dixie Runner (cultivar): dormancy and cold storage, 646; pod number and yield variation, 432; and yield increases, 104; yield stability of, 105-106
- Dixie Spanish (cultivar): amino acid composition, 658T; fatty acid composition, 661; and herbicide injury, 241; physiological maturity and, 632-633; resistance of, 300
- DMI, 628-629
- DNA transcription, 646
- Dogs, and toxicity testing, 229-230
- Dollar value, and maturity, 624
- Dolomite, as liming material, 141
- Domestic market, 748
- Don José Campos, 14
- Dormancy, 65, 102, 416-417, 635, 641; after-ripening effects, 643-644; rate of curing and, 642; defined, 642; developmental aspects, 642-643; and postharvest chemical treatments, 644-646; and storage effects, 646-647
- Double acid extractant: composition, 152; critical level, 152
- Dough conditioners, 726, 728
- Doughnuts, peanuts used for, 729
- Down windrows, 468
- Drainage, 242
- Driedfruit beetle, 522

- Drill spacing, 134
 Drought, 85, 113, 136, 466, 492; effect on apparent photosynthesis, 423; avoidance, 178; effect on calcium uptake, 172-173, 413; effect on photosynthesis, 422-423; effect on plant processes, 165-183; effect on pod yields, 430; research needs, 757-758, 760-761, 763; sensitivity, 182; relative tolerance to, 43, 63, 164, 180-181
 Drought stress, 180; and aflatoxin accumulation, 499-500, 513; and AMI, 631-632; and leaf water potential, 190; and maturation process, 627; and SHMI, 629
 Drugstore beetle, 540F; type of damage, 539; description, 537, 539; life history, 539
 Drum drying, 702, 709, 717, 719, 724
 Dry-basis moisture content, 461-462
 Dry heat, and trypsin inhibitor, 680
 Drying, 487; costs of, 482; vs. "curing," 475; and insect infestation, 553; and moisture content, 469; and mycotoxin invasion, 490-492, 498; rainfall effects, 469-470; temperature effects, 468-469; uniformity of, 479; rate in windrow, 467-468. *See also* Artificial drying, Curing
 Drying bins, 478-479
 Drying parameter, 475; defined, 468; and moisture content, 469
 Drying systems: alternative, 483-484; bin or wagon, 478-479; single trailer, 582
 Drying temperatures: effects on disease invasion, 378; on flavor, 477; on insect infestation, 553; on milling quality, 476
 Drying wagons, 473, 478-479, 743; cut-away view, 478F; and heat requirements, 481; and insect infestation, 553
 Drying zone, 478F, 479
 Dry matter accumulation: and evapotranspiration, 190; inheritance of, 76; as a measure of maturity, 631; and reproductive growth stages, 181; and soil water deficiency, 168; effect of temperature, 430
 Dry matter growth rates, 427-428
 Dry roasting, 656
 Dry rot, 368
 DTPA-TEA, 156
 Dual, 225T
 Dump pit, 572, 579, 581F, 607
 Dust insecticides, 284, 288. *See also* Insecticides
 Dutertre, Jean Baptiste, 4-5, 9, 11, 14
 Dwarfism, genetics of, 55-56
 Dymid, 225T
 Dynamic pressure, 579

E

- Early Bunch (cultivar): characteristics of, 130-131; early maturity, 114; require-

- ments of, 131; resistance of, 280; yield stability, 106; and yield increases, 104
 Early Hegari (sorghum), 389
 Early history, of peanut, 1-12
 Early leafspot, 332, 350F; control of, 329; sporulation, 327. *See also* Leafspot diseases
 Early maturity, 108, 114, 117
 Early nodulation, 180
 Early ovule abortion, 412
 Early planting. *See* Planting time
 Early Runner (cultivar): amino acid composition, 658T; aflatoxin production, 497, 501-502; correlation coefficients, 75-76; cytoplasmic inheritance, 86; dormancy and cold storage, 646; response to herbicides, 241; heritability estimates, 75-76; and nitrogen fertilization, 148; resistance of, 111, 300, 388; and yield increases, 104; and yield stability, 105
 Earthnuts peanut synonym, 10
 EC-20979, 288
 EC 76446(292), 340
 Ecological shifts, in weed populations, 243
 Ecology of pests, 520-521, 523 et passim.
 Economic thresholds, 308; concept of, 309; defined, 257; and insecticide applications, 280, 298
 Economics: and irrigation, 164; and optimum water use for yield, 197
 Ecosystem, as management unit, 251
 Edaphic conditions: and germination quality, 418; and soil-inhabiting pests, 292-293
 EDB, 389
 Edible export consumption, 742
 Edible market seed: milling diagrams, 605F, 606F; operational flow chart, 591F; shelling operation, 591F; as primary use of seed, 590, 592
 Edible protein products, 721-731; and production technology, 701
 Effective Langley's Index, 625-626
 Effective rooting depth, 193
 Egg white, coagulated, 719
 Egret (cultivar), 100
 Eicoseonic acid, 661T, 662T, 663T; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T
 8-hydroxyquinoline, 38-39
 Elaterids, 301
 Electrical resistance blocks, 197-198
 Electrolyte leakage, 644
 Electronic color sorters, 603
 Electronic counter, 569T
 Electronic moisture tester, 587
 Electronic sorting, 618
 Electronic sorting machines (ESM), 745
 ELK. *See* Extra large kernels
 Embryo culture, 447, 449. *See also* Tissue culture

- Emulsification, 713, 715-717
 Emulsifying properties, 715
 Emulsion capacity, 716
 Emulsion spray, 555
 Emulsion viscosities, 715
 Enclosed screw conveyor, 579
 Endogenous hormone levels, 417
 Endogenous metabolic barriers, 417
 Endogeocarpic mycoflora, 489, 505
 Endoparasites, 382
 Energy balance considerations, 187-188
 Energy budget techniques, 184
 Energy: conservation, 201, 761; consumption, 473, 478, 481-482; cost estimates, 482; requirements, 475, 482; shortages, 757; utilization, 761
 Energy value, of peanut seed, 679
 Enide, 225T
 Environment: effects on growth, 430-431, 437-440; effects on nitrogen fixation, 445-446
 Environment x genotype interactions. *See* Genotype x environment interactions
 Environmental conditions: and aflatoxin invasion, 494-496, 504; effect on evapotranspiration, 189-191; and harvest time, 629; interaction with cultivars, 658; and maturation process, 627; and pod damage, 487; viscosity of protein molecules, 713; research needs, 762-763; and weed control, 242
 Environmental conditions affecting herbicides: soil fertility levels, 237; soil moisture, 233-236; soil organic matter content, 238; soil pH, 237-238; soil type, 238-239; temperature, 236-237
 Environmental correlation coefficients, 104
 Environmental demand and crop evapotranspiration, 189
 Environmental factors affecting nitrogen fixation: light, 445-446; temperature, 446; water stress, 446
 Environmental factors regulating plant growth, 430-431; irradiance and quality, 437-440; light, 437; photoperiod, 440-441; temperature, 441-443
 Environmental Protection Agency (EPA), 224; and fumigants, 391-392; label requirements, 229, 231; registration requirements for herbicides, 228-229; ruling on use of Kylar, 135
 Enzymatic activity, 413
 Enzymes, 633, 636; hydrolysis, 710, 721
 Epistasis, genetic variance, 81-82
 Epistatic gene interaction, 99
 EPTC, 222-223; control of annual weeds, 222; control of stem rot, 353; subsurface application, 223
 Equilibrium moisture content, 463-466; bin or wagon drying, 478-479; and rate of drying in windrow, 467; Henderson equation, 465; effect of reduced relative humidity, 476; Smith equation, 465
 Equipment arrangement, 603
 Equipment management, 625
 Erect growth habit, 52; and gibberellin activity, 433
 Ergotism, 486
 Erucic acid, 665T
 Erythromycin, 70
 Essential amino acids, 679; contents in peanuts, 682T; nutritional quality of peanut protein, 680; exceeding recommended daily allowances, 681; requirements of man, 682. *See also* Amino acids
 Essential fatty acids, 682. *See* Fatty acids
 Esters, 677T
 Esters styrene, 676T
 Ethanol, 636-637, 640-641
 Ethanol-insoluble materials, 668T
 Ethanol-soluble materials, 668T
 Ethephon, 434, 644; and germination, 417, 645
 Ethidium bromide (EB), 70, 97, 103
 Ethnological comparisons, 15-16
 Ethoprop, 304, 391-392; control of stem rot, 353
 Ehtrel, 438, 645-646; inhibition of flowering, 434, 435F; stimulation of germination, 417; and yield increases, 436
 Ethyl acetate, 640
 Ethylene, 644; dormancy regulation, 417; and germination, 412, 416, 645-646; growth regulation, 433; and pegging, 434, 436; role of, 417; storage effects, 649-650
 Ethylenediaminedi-(o-hydroxyphenylacetate), 156
 Ethylene dibromide, 389-391
 Ethylene gas, 417
 Ethylene production: effect of irradiance, 438; effect of season and location, 412; effect of storage and relative humidity, 414; effect of subfreezing temperatures, 413
 Ethylene-synthesis capacity, 643-644, 646
 Ethyl 4-(methylthio-*m*-tolyl-isopropylphosphoramidate, 391
 Ethylmethanesulphonate (EMA), 70-71, 103
 Etymology, 9, 10, 16
 Evaporation, 191; defined, 183; energy balance considerations, 187
 Evaporative demand (E_0), 178; defined, 190; effect of climatic conditions, 189-191
 Evaporative energy flux (LE), 188, 191

Evapotranspiration (E_t), 181, 183-193; characterization of, 187-189; effects of climatic factors, 189-191; effect of cultural practices, 192-193; defined, 183-184; and drought avoidance, 178; and irrigation strategy, 195; ground cover and plant effects, 191; seasonal trends, 184-185; effect of seed growth and maturation, 183, effect of soil water deficit, 192. *See also* Evaporation, Irrigation, Moisture, Photosynthesis, Soil water deficit, Transpiration, Water use efficiency

Excrement damage, 533, 535, 537, 539, 544, 547, 549

Exocarp, 628, 634

Exotic 5 (cultivar), 291

Expeller, 702-703

Expeller cake, 703

Exportation/export channels: contracting 742; largest user countries, 750; European users, 747; international market, 748-750; role of U. S. government, 738

Extension services, future challenges of, 764

Extension specialists, future role of, 764

Extractable water, 192

Extraction, 689-695; methods of, 721

Extra large kernels (ELK), 612; effect of shading, 438; and water deficit, 173-174

Extruded foods, 730

Extrusion processing, 730

Eye effects, toxicity of pesticides, 229T

F

F 334A-B-14, 109, 659T

F392-8 131

F420-100, 371

F439-16-6, 662T, 664T

Facility cleanup, in insect control, 553

Fading time, inheritance of, 58

Fair Average Quality (FAQ) groundnuts, 750

Facility treatments: residual insecticides, 554; sanitation, 553-554

Facultative parasites, 487

Fall armyworm, 276-277; resistance to, 111-112

Fall panicum. *See* Panicum, fall

False wireworms, 301

Fancy size, 587, 599

Fans, 480-481, 564, 583, 745

Farm bill (1981), 741-742

Farm Bill of 1982, 747

Farmers stock markets, 747

Farmers stock peanuts: bulk density of, 579, 580T; cleaning, 572-574; curing operations, 458, 482; defined, 738; foreign materials in, 572, 573T, 574T; fumigation treatments, 559-561; grading proce-

dures, 580, 571, 586-590; insect damage and control, 520, 543, 552-554, 556; insecticide resistance, 564-565; mycofloral invasion, 493, 499; precleaning, 607; sampling, 586; storage, 557, 574, 602, 647; and support prices, 752. *See also* Peanuts, Seed stock

Fatty acid composition, 633; assessment of, 115; effect of herbicides and insecticides, 240; inheritance of, 76; of oil, 660, 661T-664T; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T

Fatty acid oxidation, 510

Fatty acid profile, 634

Fatty acids, 661T-664T

Fecal pellets, 528, 529, 537, 539, 541, 546

Feces damage, 533, 535

Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), 224; 1972 Amendment to, 231-232, 245

Federal-State Inspection Service (FSIS): aflatoxin inspection, 508, 598-599; grading procedures, 596, 598; sampling, 744

Federation of Oil, Seed, and Fats Association (FOSFA), 746

Fe-EDDHA, 156

Feedlot manure, 156, 158

Fensulfothion, 391-392; control of stem rot, 353

Fentin hydroxide, 279; control of spider mites, 330

Fermentation processes, 717

Fermented peanut milk, 722-723

Ferrous ammonium sulfate, 156

Ferrous iron, 156

Ferrous sulfate, 156

Fertility of peanut, 756. *See also* Dormancy, Germinability, Germination

Fertility, soil: effect on evapotranspiration, 192-193; effect on herbicides, 231, 237; and weed control, 242-243

Fertilization, 139-158; and aflatoxin resistance, 505-506; broadcast application of, 128; direct application, 103, 128, 756; influence on quality, 412; and irrigation, 201; micronutrients, 129; and nitrogen fixation bacteria, 128; research needs, 756-757; resistance to, 112; soil tests, 128; time of application, 128; and weed control, 242

Fertilizers: boron, 154-155; calcium, 141-146; cobalt, 155, copper, 155-156; iron, 156-157; lime, 139-141; magnesium, 147-148; manganese, 157; molybdenum, 157; nitrogen, 148-151; peanut meal, 590; phosphorus, 151-153; potassium, 146-147; sulfur, 153-154; zinc, 158

Fertilizer utilization, 125

Ferulic acid, 643

FESR-5-1-B-b4, 107

FESR-5-2-B-b4, 107

Fiber, crude, 705T

Field capacity, 137

Field curing/drying, 466-470; losses, 470; effect of moisture content, 469; rainfall effects, 469-470; rate of drying in windrow, 467-468; temperature effects, 468-469. *See also* Curing, Drying

Field drying cycle, 498

Field fungi: characters of, 487-488; during curing process, 490-492; before maturity and digging, 488-489; at maturity and digging, 488-489; mycoflora of soils, 488. *See also* Fungi

Field losses, 459, 470-471, 505-507. *See also* Yield losses

Field performance, 418

Field preparation, 458

Filtration methods, 706

Fish, and dinoseb toxicity, 226

Fish meal, 486

5-BU, 70

5-hydroxy-4-nonenic acid, 673

5,6-dihydro-2-methyl-1,4-oxathiin-3-carboxanilide, 352-353

Flakes, peanuts used for, 702, 725

Flat cultivation, 127, 217-218

Flat grain beetle, 538F; type of damage, 537; description of, 535; life history, 535, 537

Flat screeners, 614-616

Flat steel bar commercial sheller, 609, 611T

Flat-type storage, 576, 578T

Flatulence sugars, 705

Flavone (yellow pigment), 57

Flavonoids, 29

Flavor, 635-636, 671; and "aging period," 640; CLER flavor score method, 678; enzymic origin, 637-639; "green" or "beany," 636-637, 707; and growth regulators, 756; characteristics of lipoxigenase, 639-640; and monosaccharides, 671-672; volatile compounds, 640-641, 673. *See also* Aroma

Flavor change/loss: due to curing process, 458; 482-483; due to fumigation, 560; and infrared drying, 483; effect of malathion, 554; from processing for flour, 703-704; and quality loss, 574; temperature effects, 477

Flies, 522-523

Flooding, 206, 235

Floral characters: albinism, 59; chlorophyll deficiencies, 59-61; corolla color, 57; fading time, 58; other foliage characteristics, 63; inheritance of, 57-63; leaf color, 58-59; leaf shape, 62-63; standard crescents, 57-58; stem pigmentation, 63; stem

pubescence, 63; sterility, 58; wing-petal shape, 58

Floral initiation. *See* Flowering

Florida beggarweed. *See* Beggarweed, Florida

Florida pusley. *See* Pusley, Florida

Florida Jumbo (cultivar), 659T

Florida 393 (cultivar), 131

Florida 393-7-1 (cultivar), 661

Florigiant, 99, 130, 391, 459, 470; aflatoxin production, 501; and AMI, 631-632, 667T; bulk density of, 580T; calcium concentration and germinability, 143F, 144; calcium content, 667T; ambient calcium requirement, 144; and DMI, 629-630; drought, water use and E_t, 173-174, 182, 185, 186F, 196-197; fatty acid composition, 115, 661, 662T; free sugar content, 667T; fruiting habit, 130; growth habit, 130; maturity of, 114; milling quality, 592; mineral element content, 669T; oleate/linoleate ratios, 664T; pod thickness distribution, 594-595; stability of, 84, 105-106; susceptibility/resistance, 71-72, 106, 108-109, 111, 280, 300, 356, 358, 371, 387-388

Floripan, 132; collar rot resistance, 108-109; yield stability, 105

Floripan runner: hybrid vigor, 78; and oil content, 114

Florunner, 99, 117, 123, 132, 276, 391, 432, 627, 646, 680; aflatoxin tolerance, 110; and AMI, 631-632; bulk density of, 580T; and calcium, 142-144, 146; and DMI, 629-630; drought, water use and E_t, 169-170, 173-174, 175F, 177F, 179, 185, 186F, 196-197; dry matter accumulation, 181F, 428F; fatty acid composition, 115, 661, 662T; magnesium content, 147-148; maturity, 114; milling quality, 592; oleate/linoleate ratios, 664T; photosynthesis/apparent photosynthesis, 421, 423, 424F, 426F, 427F; pod maturity profile, 628; pod thickness distribution, 594-595; resistance of, 107-108, 110-111, 113, 300, 329, 334, 388; root growth, 165, 166F; row spacing, 134; seedcoat splitting, 116; and yields, 104, 134, 139, 239-240; yield stability, 84, 105-106, 108, 132

Flotation-sieving method, 385

Flours, peanut, 702-704. *See also* Defatted peanut flour

Flowering: and calcium levels, 142-144, 756; effect of cultivation, 135; and growth regulators, 434, 756; effect of high humidity, 441; initiation of, 170, 429; effect of planting conditions, 133; thrips damage and rate of, 287; effect of water deficit, 169-170

- Flowers: characteristics, 57-63; fading time, 58; and foliage consumers, 276; production of, 256
- Flow meter, 555
- Flow pipes, 607
- Flow rates, 581; and gravity table, 618
- Fluometuron, 353
- Fluorescence, 508. *See also* Inluorescence
- Flux density of water, 184
- FMI, 628-629
- Foam formation, 717-719
- Foliage blight, 362-363
- Foliage calcium, 146
- Foliage characteristics. *See* Floral characters
- Foliage consumers 253, 254F, 255, 270; flower consumption, 276; intracellular feeders, 281-292; lepidoptera, 271-281; rate of, 276-277; worldwide pest status, 274, 309
- Foliage-inhabiting pests, 271; foliage consumers, 271-281; intracellular feeders, 281-292
- Foliage uses, 474
- Foliar diseases: alternaria leafspot, 337; anthracnose, 335-336; cercospora and cercosporidium leafspots, 326-330; choanephora leafspot, 339; cowpea mild mottle, 345-346; groundnut crinkle, 346; groundnut eyespot, 346; groundnut leaf blight, 337; groundnut rosette, 340-341; marginal chlorosis, 347; melanosis, 338; peanut chlorotic leaf streak, 347-348; peanut clump, 344-345; peanut green mosaic, 346; peanut mottle, 339-340; peanut stunt, 343-344; peanut yellow mottle, 347; pepper spot and leaf scorch, 334-335; pestalotiopsis leafspot, 338-339; phomopsis leaf and stem diseases, 337-338; phyllosticta leafspot, 338; powdery mildew, 338; Rhizoctonia foliar blight, 337; rugose leaf curl, 347; rust, 330-332; scab, 336; tomato spotted wilt, 341-343; web blotch, 332-334; zonate leafspot, 337. *See also* names of individual diseases
- Folic acid, 447, 670T, 684
- Food and Drug Administration: guidelines for wholesomeness, 678-679; inspection of market products, 509
- Forced draft ventilation, 584-585
- Foreign Agricultural Service (FAS), 740
- Foreign grain beetle, 546F; type of damage, 547; description, 546; life history, 546-547
- Foreign material: and "bridging," 578; increase in bulk density, 579; combining and, 472; damage to equipment, 603; damage to seed, 592; and edible stock, 607; and gravity table, 616; handpicking, 603, 618; quantities and types of, 572-574; 597T, 599T; in raw and roasted peanuts, 678; reduction methods, 575T; source of excess moisture, 582
- Formaldehyde, 511
- 4-chlorobenzyl-tri-n-butylammonium bromide, 434
- 4-(methylsulfonyl)-2,6-dinitro-N-N-di-propylaniline, 222
- 4-(2,4-dichlorophenoxy) butyric acid, 223
- Foxtail, 208T, 232T *See also* specific names of foxtails
- Fractionation, 639, 689 et passim.
- Frankfurters, peanuts used for, 725
- Frass, 524, 526F, 528, 529, 533, 535, 537, 539, 543, 544
- Free arginine, 630-631
- Free enterprise system, 764
- Free fatty acids (FFA), 633-634, 647, 705T; carbonyls, 641; and curing, 636; and fungal growth, 496; and germination, 413; in oilstock peanuts, 660T; 751; effect of red flour beetle, 530. *See also* Fatty acids
- Free sugars, 667T
- Freeze drying, 709, 716-717, 719
- French bean, 344, 346
- Fresh weight seed/hull maturity index (FMI), 628-629
- Frost probability, and digging time, 459
- Frozen desserts, peanuts used for, 723
- Frozen fertilizer, 128
- Fructose, 666-668, 669T
- Fruit, 5,6,487; rate of addition, 104; breeding objectives, 103-104; of cultivars, 130, 132; effect of drought, 169; moisture requirements, 181; effect of planting conditions, 133; and plant turgor pressure, 183; production, 256; effect of rainfall, 125; soil mineral content and development of, 139, 144-146. *See also* Pericarp, Pod
- Fruiting zone: and excess phosphorus, 145, 412; and soil calcium level, 144, 412, and soil water, 171
- Fruit size: additive and nonadditive genetic effects, 80; as breeding objective, 104; broad-sense heritability estimate, 76; combining ability estimates, 79-80; genotypic correlations, 76-77; heritabilities, 76-77; phenotypic correlations, 76-77, 105. *See also* Pod size
- Fruit yield: Combining ability, 80; correlation coefficients, 77; heterosis, 78; hybrid vigor, 78. *See also* Yield, Yields
- Full season irrigation, 136
- Fumigant nematocides, 389-391
- Fumigants: conventional, 559-561; modified atmosphere, 561

- Fumigation, 583, 585
- Functional properties: emulsification, 713, 715-717; foam formation and stabilization, 717-719; gelation, 719-720; oil absorption, 721; solubility, 707-712; viscosity, 713; water sorption, 720-721
- Fungi, 486; combining ability for resistance to, 80; control of, 505-511; field, 487-492; factors affecting growth, 497-505; other types of, 511-513; parasites of, 520; pod and seed damage, 116, 503-504; production of, 496-497; as enemy of spider mites, 290; storage, 492-496
- Fungicides: methods of application, 329-330, 414; control of *A. flavus*, 507-508, 513; in combination with copper, 155; control of disease, 335, 356-359; 373-375, 378; effect on germination, 648; and management of leafspot, 136, 327, 335; effect on maturation, 631; control of nematodes, 387; organic mercurial, 367-368; research needs, 761; treatment of seed stock, 603, 618-619; and spider mite outbreaks, 258-259. *See also* names and chemical formulae of individual fungicides.
- Furans, 672-673, 675T-677T
- Fusarium diseases, 364-365; causal organism, 365; control of, 366; disease cycle and epidemiology, 365-366; symptoms of, 365
- Fusarium wilt, 365
- G**
- GA. *See* Gibberellic acid, Gibberellins
- Ga. 61-42, 658T
- GA 119-20, 131
- Galactose, 667-668
- Galacturonic acid, 668T
- Gamma-ray mutation, 97; to induce cytoplasmic mutations, 86; to increase drought resistance, 113; rate of, 71
- Gamma transmission instruments, 198
- Gas chromatographic technique, 657, 673
- Gear pump, 555, 557
- Gel electrophoresis, 656, 690, 693
- Gelation, 719-720, 723
- Gelation gels, 719
- Gelechiids, 273
- Gene action, 754; combining ability, 79-80; epistasis, 81-82; heterosis, 78-79; type of, 72-73, 77-82; variance studies, 80-81
- Gene pools: for aflatoxin resistance, 506-507; for improvement of peanut, 42, 45, 95-96. *See also* Germplasm
- General combining ability (GCA), 99-100; additive genetic effects, 79, 100; and disease resistance, 80; importance of, 79; and yield stability, 104
- Generation means analysis, 80-81
- Genetic advance, 74
- Genetic bases: inheritance of albinism, 60T; inheritance of growth habit, 54T; testa color inheritance, 66T-67T
- Genetic behavior, 105
- Genetic correlations: scarcity of, 85; of yield, 103-105
- Genetic control, 433
- Genetic divergence: in *Arachis*, 40; as a result of mutations, 97
- Genetic improvement, 42-44, 95 et passim.
- Genetic linkage, plant characters and disease, 72
- Genetics, 50-87; breeding goals, 95, 103-117; breeding strategies, 41-45; breeding systems, 99-103; cultivar maintenance, 116-117; cytoplasmic inheritance, 85-87; effects on growth, 431; inheritance of qualitative characteristics, 52-72; inheritance of quantitative characteristics, 72-85; research needs, 754-755
- Genetic variability/variance, 50-52; in breeding systems, 99; in genotype x environment interactions, 82; of macromutant "cup," 103; nature of, 72-73, 75; in plasmone, 97; quantitative traits, 80-81; sources of, 95-99
- Genomic divergence, 40; homologies, 36, 39-40, 44
- Genotype x environment interactions, 82-83; evaluation of stability, 83; effect on natural cross-pollination, 96-97; effect of time of year, 442; effect of water management systems, 443; and yield stability, 105-106
- Genotype x pesticide interactions, 113
- Genotype stability, 83-84
- Genotypes, chlorophyll-deficient mutants, 61T
- Genotypic coefficient of variability, 74-75
- Genotypic correlations: and early maturity, 114; and heritability, 65; vs. phenotypic correlations, 75; for yield, fruit size and maturity, 76-77, 104
- Genotypic variance, 74, 180-181
- Geocarp, 486-488
- Geocarposphere, 367, 488, 504-505
- Geographic distribution: of *Arachis*, 26F, 27F, 50-52; and genetic variability, 51F; of peanut production in U. S., 123
- Geographic location, effect on seed quality, 412
- Geographic origin, of peanuts, 12-13

- Georgia 186-28, 388
 Georgia Station runner, 304
 Germinability, effect of water deficit, 174
 Germination: after-ripening effects, 643; in calcium-deficient plants, 143-144; effect of drought, 174; and ethylene production, 644-646, 760; and fumigation, 560-561; and fungicide treatment, 648; influence of harvesting, curing and handling conditions, 413-414, 458, 492; and imbibition, 414-415; and infrared drying, 483; effect of insect pests, 530; effect of land preparation, 126; metabolism, 415-416; effect of preharvest conditions, 412-413; production management, 462; quality evaluation, 418; and refrigeration, 648; seed dormancy, 416-417; seedling development, 419; effects of storage, 414, 494-496, 647-650; temperature effects, 413, 415
 Germination inhibitor concentrations, 642
 Germination tests, 618
 Germplasm accessibility, 42
 Germplasm, peanut: collections of, 95-96, 754; diversity, 85; drought tolerance, 180-181; fatty acid composition, 662, 664; maintenance of, 98-99, 754; research needs, 759; storage of collections, 98-99
 Germplasm pool, diversity in, 73-75
 Germplasm, wild species, 44, 755
 GH 119-20, 85
 GK-3, 108
 GKP 10017, 72
 GKP 10602, 330
 Giant foxtail, 207. *See also* Foxtail
 Gibberellic acid (GA): and enzyme activity, 416-417; and germination, 416-417, 645; growth factor, 53, 56
 Gibberellins, 433-434, 643, 646; culture of reproductive tissue, 449; and dark requirement for pod development, 439; effect of light and, 438; and peg growth, 434
 Glass, as foreign material, 573T, 575T, 618
 Globulins, 689-690
 Glucose, 666-668, 722; favorable effect on *A. flavus*, 146; in pigments, 68; effect of roasting, 669T
 Glutamic acid, 656, 657T-659T, 700-701; and flavor, 671
 Glutamine, 671
 Glyceraldehyde, 672
 Glycine, 657T-659T, 701
 Glycolipid, 666
 Glycoproteins, 656
 Glyoxylate cycle, 415
 Glyphosate: nomenclature, 225T; relative toxicity of, 230T, technical description, 227; weeds controlled by, 232T
 Gnotobiotic conditions, 489, 504-505
 Goosegrass, 208T, 209F; response to herbicides, 225, 232T; mechanical removal of, 208; proclivity, 214, 215T, 216T, 218T
 Government regulations, 737-740, 764. *See also* USDA, U. S. government
 Grade factors, 586
 Grade procedures, 587T
 Grading of peanuts, 580-590, 744; of cleaned inshell and shelled peanuts, 595-600; research needs, 762
 Grading screens, 596T
 Grain residues, and lesser mealworm, 549
 Grain sorghum, 125-126; 372-373
 Grains, 486-487, 495, 506, 513; bacterial control of insects, 566; emigration of thrips from, 286, 288; as foreign material, 573T; and insect damage, 533, 544, 549; and warehouse designs, 579
 Granaries, 539, 541
 Granular fertilizer, 128, 145
 Granular insecticides: control of *A. cracivora*, 284; of *E. lignosellus*, 298; of grubs, 307; of *Tetranychus*, 291; of thrips, 288
 Granulosis virus, 566
 Grasses: as a crop, 504, 506; as a pest, 214, 227, 573T, 575T
 Grates: designs of, 610T-612T; in shelling process, 609
 Gravimetric dig, dry, and weigh method, 198
 Gravity tables, 616-618
 "Gray mold," 374
 Green foxtail, 207. *See also* Foxtail
 Green harvesting method, 116
 Green peach aphid, 344
 Green rosette, 340
 Greenville sandy clay loam, 238-239
 Grits, peanuts used for, 702-704
 Groundbean, 327
 Ground beef patties, peanuts used in, 724-725
 Ground cover, effects on evapotranspiration, 191
 Ground cracking, 135
 Groundnut, 486, 488; and moisture equilibrium, 459T. *See also* Peanut
 Groundnut crinkle, 346; causal agent, 346
 Groundnut crinkle virus (GCV), 346
 Groundnut eyespot, 346
 Groundnut eyespot virus, 282
 Groundnut leaf blight, 337
 Groundnut root rot, 368
 Groundnut rosette virus (GRV), 282T; aphid vector, 281-282; causal agents, 341; control of, 341; control of vectors, 285; epidemiology, 341; resistance to, 71; secondary spread, 283; symptoms of, 340-341; types of, 284
 Growers, peanut: contracting, 741-743; delivery and sampling, 743-744; harvesting, 743; laws and regulations governing, 737-741; marketing risks and decisions, 737, 742, 751; needs of, 95, 117, 756; effect of supply and demand on, 747
 Growing techniques, role of U. S. government, 738
 Growth analysis: dry matter growth rates, 427-428; environmental effects, 430-431; reproductive growth, 429, 430; vegetative growth, 428-429; yield variation, 431-433
 Growth duration, 755
 Growth habit: of cultivars, 131-139; and environmental factors, 53, 430-433; inheritance of, 54T, 55; types of, 52
 Growth physiology, 411, 449-450; environmental factors, 437-443; germination and seedling growth, 411-419; growth regulators, 433-437; nitrogen fixation, 443-447; photosynthesis and growth analysis, 419-433; research needs, 760; tissue culture, 447-449
 Growth rate, 428; of dry matter, 428; and pod removal, 255
 Growth regulators: and drought stress, 136; and Kylar, 135-136; injection by spider mites, 290; pod and seed formation and, 435-437; reproductive growth, 434-435; research needs, 756; vegetative growth, 135, 433-434
 Growth retardants: effects of ABA, 434; CCC, 433; morphactins, 433-434; SADH, 433; yield enhancement and irrigation, 168
 Gruels, peanuts used for, 730-731
 Guayuru (cultivar), 336
 Guazatine triacetate (GTA), 280
 Guinea ant, 523
 Geunea pigs, 512
 Gujarat Dwarf (cultivar), 56
 Gujarat narrow leaf mutant, 62-63
 Gypsum: as calcium source, 129, 145; in disease control, 362; elimination of plumule damage, 413; research needs, 757; as source for sulfur, 153; as used in tensiometers, 198; "wet" 145, yield response, 139-141, 146

H

- Haemoglobin content, 151
 Hairy fungus beetle, 548F; type of damage, 546; description of, 544; life history, 546
 Hairy indigo, 208T, 216T, 218T
 Hairy vetch, 333
 Half cystine, 656, 657T-659T, 679, 700-701

- Handling, 574-586, 607-608; influence on germination, 413-414; new methods of, 601-602
 Handling Committee, 601
 Hand sorting, 509, 603, 618
 Hand weeding, 213
 Harvest conditions, and germinability, 413
 Harvesting, 743; and aflatoxin contamination, 507-508, 513; damage during, 487; defined, 458; mechanics of, 460-461; pre-digging operations, 460; research needs, 760-761; timing of digging, 458-460
 Harvesting techniques, 116; effect of early drought, 171; and insect infestation, 552; and milling quality, 593; research needs, 761; effect of weeds, 214, 244
 Haulm, 487
 Hay, peanut foliage used for, 474
 Hazard indicators, of pesticide toxicity, 229T
 Heater capacity, 481
 Heat pump drying systems, 483
 Heat requirements, during curing, 481
 Heat stress, 760
 Heat unit system, 625
 Hedging techniques, 751-753
 Hemicelluloses, 668
 Henderson equation, 465-466
 Heptachlor, 303
 Heptadecenoic acid, 661T
 Heptadecylic acid, 661T, 665T
 Herbicide—crop rotation concept, 244-245, 758
 Herbicides, 134, 241-242; methods, time and rate of application, 135, 220-228; in combination, 221-222, 239-241; external effects on activity and persistence, 231-239; and irrigation, 164, 201; labeling requirements, 231; effect on market quality, 221-222; effect on nitrogen fixation, 150; and pest control, 352, 357; research and development, 219-224, 758; technical descriptions, 224, 225T, 226-228; relative toxicity, 226T; toxicology, 228-231; and weed control, 135, 219, 231, 232T, 242, 244-245. *See also* individual names and formulae of herbicides.
 Heritability: in breeding systems, 99; of iodine value, 115; for pod size and yield, 77; and resistance, 110; of seedcoat splitting, 116; in segregating generations, 75-77; for yield, fruit size and maturity, 76-77, 114
 Heritability values, 74
 Heterosis: gene action, 78-79; and yield stability, 105
 Hexanal, 636-639, 641, 673
 Hexane-acetone-water extraction, 510
 Hexane-ethanol-water extraction, 510

Hexane extraction, 702-703
 Hexane-methanol extraction, 510
 Hexane residue, 703
 HG-4, 668T
 HG-10, 288, 291
 Hide beetle, 521
 Hip-rip treatments, 352
 Hirst spore trap, 334
 Histidine, 657T-659T, 671, 701
 History of peanuts: archaeological evidence, 11-12; chronological, 1-11; geographical evidence, 12-16; industrial developments, 16-18; research needs, 754. *See also* Origins
 Hoeing, in weed control, 240
 "Hollow heart," 129, 154, 172
 Homogenates, 638
 Honeyvine milkweed, 208T; response to herbicides, 232T; proclivity, 216T, 218T
 Horsenettle, 208T, 244; response to herbicides, 232T; proclivity, 214, 215T, 216T, 218T; research needs, 758
 Host-bacteria symbiosis, 444
 Host resistance, 759
 House fly, 522
 Hull processing equipment, 607-608
 Hulls: internal color, 626, 632; shelling and thickness of, 594-595; uses of, 608. *See also* Pericarp
 Humidistat, 477
 Humidistat-controlled aeration fans, 745
 Humidity, 187-188; and cumulative pan evaporation, 200; and deterioration of seed, 650; effect on evaporative demand, 189; effect on flowering, 170; and growth of fungi, 487, 494-500; and net photosynthesis, 176; during storage, 648-649. *See also* Moisture, Relative humidity
 Hybridization: artificial techniques, 100-101; in breeding systems, 99-101; genetic consequences, 75-77; genetic variability, 50-52; heritability estimates, 75. *See also* Genetics
 Hybrid progeny management: branching pattern variations, 101; plant spacing, 101
 Hybrids: artificial, 97; behavior of, 42; coding problems, 98; and drought tolerance, 180-181; natural, 96; persistence of vigor, 78, 99-100. *See also* Cultivars
 Hydration, 414-415
 Hydraulic lift, 579
 Hydraulic pressing, 702
 Hydrocarbons, 676T-677T
 Hydrogen peroxide, 510-511
 Hydroperoxyoctadecadienoic acid, 639-640
 Hydroquinone, 638
 Hydroxylamine (HA), 70

Hygroscopic equilibrium moisture content, 582, 583F
 Hysteresis, 463

I

IAA. *See* Auxins
 Ibimani (peanut synonym), 9
 ICR-170, 70
 Identification methodology, 746
 Illite clay, 238
 Imbibition, 414-415
 Immature class, 626, 628, 635-636
 Immature pods, removal of, 575T
 Immature seed: colonization by *A. flavus*, 489-501, 503, 505; germinability, 413; off-flavor, 477
 Imine, 672
 Immuno-electrophoresis, 29, 692
 Impact damage, 413
 Importation, 750, 752
 Imports, 16
 Improved Virginia Bunch, 185
 Inbreeding, genetic consequences of, 75-77
 Inbreeding depression, 78
 Incas, and peanut cultivation, 1, 7, 9
 Incompatibility barriers, 755
 Indemnification program, 739
 Indian meal moth, 525F, 526F; bacterial control of, 566; type of damage, 524; description of, 523-524; life history, 524; predators of, 551; resistance of, 565
 Induced drought, 176
 Industrial development, 16-18
 Industry, needs and desires, 95, 117
 Infantile liver cirrhosis, 486
 Infiltration data, 193-195
 Infiltration rates, 193
 Inflorescence, 50-51; of main stem, 55. *See also* Florescence
 Inflorescence consumption, 276
 Infrared drying, 483
 Infrared spectrophotometric method, 673
 Inhalation LC₅₀, 229T
 Inheritance: floral characters, 57-63; growth habit, 52-53, 54T; mono and digenic models, 52; qualitative characters, 52-72. *See also* Genetics, Heritability
 Inhibitory phenolic acids, 643, 650
 Initial moisture content, 481-482. *See* Moisture content
 Initial seed quality, 649
 Injury site-recovery ability relationship, 255. *See also* Spatial tolerance
 Inner testa color, inheritance of, 69
 Inoculation, 149-150; forms of inoculant, 128
 Inositol, 447, 667T, 669T-670T

Input solar energy (R_s), 188-189
 Insect control: Bacterial control, 566; integrated pest management, 250, 253; post-harvest pests, 552-566; research needs, 758-759; systems of, 580T, 584
 Insect detection procedures, 762. *See also* Sampling, Detection
 Insecticidal aerosols, 561-564
 Insecticidal prophylaxis, 280
 Insecticidal protective treatment, 554
 Insecticides: application techniques, 278, 581; and disease control, 341; and economic thresholds, 280; genotype x pesticide interaction, 113; and irrigation practices, 761-762; source of excess moisture, 582; and preharvest insects, 279-280, 284, 288, 298-300, 303-304, 307; preplant application, 252; prestorage spraying, 581; research needs, 761-762; resistance to, 279, 564-565; and spider mites, 258-259, 291; and overspace ventilation fans, 583-584
 Insect infestation: effect on milling quality, 594; quality loss during storage, 574
 Insect management, 308
 Insect pathogens, 487, 499, 505-506, 762
 Insect-transmitted diseases, 253
 Insects, 187; arthropod pests, 259-271; biology, ecology and recognition of, 520-523; damage by, 520, 523, et passim.; disease transmittal, 253; feeders on roots, pegs and pods, 253; foliage consumers, 271-281; and fungal invasion, 487, 499, 505-506; intracellular feeders, 281-292; and irrigation control, 201; postharvest, 520-552; resistance to, 111-112, 391; soil-inhabiting, 292-307; and weed control, 242. *See also* Pests, individual names of insects
 Inshell peanuts: aflatoxin analyses, 599-600; in grading procedure, 587; sampling and sorting, 595-596, 598-599; sorting for roasting, 595
 Insoluble proteins, 649
 Inspection belt, 586, 596T
 Integrated pest management (IPM), 245, 253; approaches to insect control, 251-252; conceptualization of peanut system, 252-253; crop-pest relationships, 253-257, 309; defined, 250; economic benefits of, 299-300; field sampling, 299; overall perspective, 307-310; Peanut Insect Defoliation Model, 277-278; pest status, 257-259; principles of, 251; tactical approaches, 307T

Interactions: cultivar x environment, 658; date x depth for nematode location, 386; diseases and weed control, 217-219; genotype x environment, 82-83, 442-443; genotype x pesticide, 113; of herbicides and pesticides, 239-242; between nematodes and other pathogens, 386-387; and yield stability, 105-106. *See also* Genotype x environment interactions
 Intracellular botryose haustoria, 328
 Intermediate class, 626, 628
 Internal hull color, 626
 Internal oxygen concentration, 635-636
 International Crops Research Institute for the Semi-Arid Tropics (ICRISAT), 96, 100-101, 114, 331, 345
 International market, 748-750
 Interplant movement, 270-271
 Intersectional cross-compatibility, 35-37, 40-41
 Interspecific hybridization, 25-28, 33; and aneuploidy, 30-31; artificial techniques, 100-101; improvement of peanut, 41-45; interpretation of pairing relationships, 39-40; meiosis in, 35-37
 Interspecific hybrids, 34-37, 754
 Intracellular feeders, 253, 270, 281; aphids, 281-285; spider mites, 289-292; thrips, 285-289
 Introductions. *See* Plant introduction
 Introgression, 45
 Invertase treatment, 722
 Inverted windrow drying, 468, 470, 491-492, 499, 507, 513
 "Inverter" device, 214, 460, 517
 Iodine value, 114, 660, 634; heritability of, 115
 Ion-exchange chromatography technique, 657
 Ion transfer mechanisms, 178
 Ionic environment, 708, 715, 719
 Ionic strength, 713, 720
 Iprodione, 356
 Iridescent virus, 566
 Iron: breeding for tolerance to, 757; chlorosis, 156; cultivar differences, 156-157; deficiency of, 156; importance of, 129; sources of, 156
 Iron chelate, 279
 Iron content, 444, 668, 669T
 Irradiance, 431, 437-440; response of apparent photosynthesis, 422; response of canopy AP, 426; leaf and plant age and, 424

Irrigation: daily water requirement, 136; frequency, 137, 167T; full season, 136; effect on herbicides, 135, 233; importance of, 133, 136, 164, 195, 201; monitoring of, 137; effect on nodulation, 180; overflood vs. sprinkler, 292; and pest control, 301, 356, 372, 499, 505, 761; and planting patterns, 134; research needs, 757-758, 761; scheduling, 200; seasonal water requirements, 136; influence of soil characteristics, 193-194; strategy, 195-201; surface water, 137; systems approach, 201; and water accounts, 200; effect on weeds, 213, 242; and yield reductions, 183. *See also* Water

Irrigation system, 201

Isocitric lyase, 413, 415-416

Isoelectric points, 694, 705

Isolates, 511, 702, 706-707; pathogenic, 354, 376, 388

Isoleucine, 657T-659T, 679, 701

Isopropyl m-chlorocarbamate, 220

Isopropylamine salt, in formulation for glyphosate, 227

Isotherms, 464F

Israel line, 136, 108

Ivyleaf morning glory. *See* Morning glory, ivyleaf

J

Janssen's formula, 579

Jaubert, 16

Jenkins Jumbo (cultivar), 114; amino acid content, 658T; methionine content, 115, 658T; resistance of, 111

Jimsonweed, 208T, 211F; proclivity, 214, 215T-216T, 218T

Johnsongrass, 208T, response to herbicides, 232T; proclivity, 214, 215T-216T, 218T

Jonstonus, 4, 9, 11

Jumbo Runner (cultivar), 612

'Junagadh,' 83

K

Kaolinite, 238

Kaolinite-montmorillonite, 238

Karyotypes, 32, 34, 37-38

Kernel. *See* Seed, Pod

Kernel density, 626, 628

Kernel weight, 628

Ketoamine, 672

Ketoglycerides, 640, 674T

Ketones, 636, 674T-675T, 677T

Key pest status, 257, 274. *See also* Pest status

KH-149A, 341

KH 3278, 87

Khorat soils (sandy), 153

Kinetin, 147, 433, 439, 645; development of callus, 448; in reproductive tissue culture, 449

Koboka (US 26), 506

Kojic acid, 512

Kopergaon-1, 62

Kopergaon-3, 56

Krinkle (chlorophyll-deficient mutant), 59-61, 68

Krinkle leaf mutant, 62

Krinkle lutescens (chlorophyll-deficient mutant), 59-61

Kumawu Erect (cultivar), 388

Kylar: effect on germination and vigor, 437; growth retardant, 433; effect on peg strength, 435; plant growth regulator, 135-136; effect on pod weight, 435, effect on yield, 436

Kylar 85, 135-136

L

Labat, Jean Baptiste, 4-5, 10-11, 14

Labeling, 229, 231

Laboratory sheller (Model 3), 115

Labor conservation, 201

Labor management, 625

Lactate extractant: composition, 152; critical level of, 152-153

Lactones, 676T, 677T

Lake fine sand, 165-166

Lakeland loamy sand, 141

Lambsquarter: common, 208T, 212F; proclivity, 214, 215T, 216T, 218T

Landplaster. *See* Gypsum

Land preparation: fertilization, 128-129; methods of, 126-127; objectives of, 126; and weed control, 135, 242

Land tillage. *See* Tillage

Lard, 660

Large crabgrass: proclivity, 214; and yield reductions, 213

Larger black flour beetle, 521

Larger cabinet beetle, 521, 542F; type of damage, 541; description, 541; life history, 541

Larger grain borer, 521

Larval dropping damage, 524, 529, 543

Larval parasitism, 278

Las Casas, Bartolomé, 1

Lash barges, 749

Lasso, 225T

Late leafspot, 333, 350F; control of, 329; sporulation, 327. *See also* Leafspot diseases

Latent heat of vaporization, 190

"Late-season" weeds, 207

Lauric acid, 660

Laws and regulations, 737-740. *See also* U. S. government

Leaching: reduction of dormancy, 417; of herbicides, 224-228, 237

Leaf: age and effect on AP, 424; anatomy of, 419-420; inheritance of color, 58-59; and diffusive resistance, 423; display, 178; expansion, 168, 430; gas exchange and water stress, 174-178; heat load, 178; nitrogen content and photosynthetic rate, 151; orientation, 191; and phosphorus content, 153; photosynthesis, 419-425; surface moisture and evaporative demand, 190; temperature, 178, 190; tissue content, 146-147

Leaf area growth, 256

Leaf area index (LAI): and crop evapotranspiration, 181-182, 191; defined, 425; and disease effects, 329; and drought tolerance, 180; dry matter accumulation, 428F; effect of water deficits, 168, 192; and weekly maximum canopy AP, 427

Leafhoppers, 391

Leaflet folding, 178, 419

Leaf osmotic potential (ψ), 179

Leaf relative water content (RWC): evaluation of, 179; and water stress, 174

Leaf scorch, 334, 350F; causal organism, 334; control of, 335; disease cycle and epidemiology, 334-335; symptoms of, 334, 336

Leaf shape: "cup", 62; elliptical recessive vs. elliptical oblong, 62; Gujarat narrow leaf, 62; inheritance of, 62-63; krinkle leaf mutant, 62; mottled leaf mutant, 62; spear-shaped small leaflet mutant, 62

Leafspot diseases: vs. calcium deficiency, 142; and canopy apparent photosynthesis, 427; carbon dioxide exchange and, 426; and crop rotation, 125-126; and foliar disease complex, 332; and genetics, 44-45, 71-72, 75, 77, 80, 102-103, 117; response to irrigation, 201; and Kylar 85, 136; and leaf area index, 425; resistance to 44-45, 71, 80, 102-103, 106-108, 117; and weed control, 135, 214; and yield, 108. *See also* names of individual leafspot diseases.

Leafspot fungicide, 136

Leaf thermocouple hygrometers, 179

Leaf turgor potential (ψ_p), 179

Leaf water potential (ψ_l), 177-178, 180, 422-423

Legume nodulation, 446

Lepidoptera, 271-272, 293; biology, 272-274, 293-295; damage, 274-278, 295-296; management of, 278-281, 296-301; movement, 274

Lepidopterous defoliators, 278

Lesion nematodes, 379F, 381; control of, 126, 389-393; resistant cultivars, 388; infection and spread, 382-383; resistance to, 112-113; symptoms of, 381-382. *See also* Nematodes

Lesser cornstalk borer, 111; control of, 391; and pod damage, 503, 506; pest status, 258; resistance to, 43, 111

Lesser grain borer, 521

Lesser mealworm, 550F; type damage, 549; description, 549; life history, 549

Lethalaire A-40, 563

Lethalaire A-41, 563

Lethalaire A-50, 563

Lethal concentration (LC₅₀), 230

Lethal dose (LD), 229, 512

Leucine, 657T-659T

Li-Cor 1600 Steady State Autoporometer, 176

Lifting, 487, 490-491, 498, 500. *See also* Digging

Light, 437; control of growth habit, 433; irradiance and quality, 437-440; effect on nitrogen fixation, 445-446; photoperiod, 440-441

Light intensity. *See* Irradiance

Light interception, 426-427

Light transmittance, 626-627

Lignin: antifungal properties of, 109, 507

Lignoceric acid, 662, 663T-664T; triglycerides and phosphatides, 665T

Lignosulfates, 156

Lime, 139-141; acid-soil infertility, 141; time of application, 141; as calcium source, 129; effect on herbicides, 237-238; importance to pegging zone, 139, 141; research needs, 756; yield responses, 139, 140F

Lime slurry, 141

Linkage, 72, 755

Linkage disequilibrium, 81

Linkage mapping experiments, 72

Linoleate content, 640, 673

Linoleic acid, 638-639, 660-664, 682; and maturity, 662; and oil stability, 114, 663, 664; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T

Linolenic acid, 661T, 663T; and oil stability, 665; triglycerides and phosphatides, 665T

Linseed oil, 660

Linuron, 222

Lipase, 633

Lipid content, 636

Lipids, 414-415, 429-430, 633; and curing, 636; energy sink and, 257; metabolic reserves, 415; nutritional quality, 683; radioactivity of, 636; storage of, 415; synthesis of, 636

- Lipoxidase, 636, 647
 Lipxygenase, 633, 638; characteristics of, 639-640; enzyme degradation process, 627; inactivation of, 702; reaction, 637
 Liquid cyclone process, 702, 718
 Liquid fertilizer, 128
 Liquid seed treater, 619
 Little house fly, 522
 Live and dead insect damage, 524, 528, 533, 535, 537, 539, 541, 543, 544, 546, 547, 549
 Loading equipment, 580T
 Loam soil, 193, 227-228
 Loamy sand soil, 183, 194, 197
 Logistic distribution model, 616
 Longheaded flour beetle, 536F; type of damage, 535; description of, 535; life history, 535
 Long position, 752
 Loose shelled kernels (LSK), 132, 744-745; and bulk density, 579; effect of combining, 552; and grading, 508, 587; and insect damage, 520; oil stock market and, 592; and picking cylinder speed, 472; pre-cleaning, 607; effect on storage and processing systems, 572, 574T; and windrow drying, 471
 Low-fat peanut flour, 726. *See also* Defatted peanut flour
 Low-oxygen atmosphere, 602
 Low-press expelling, 702
 Luncheon loaves, peanut used for, 725
 Luteins, 61, 671
 Lutescens 0018 (chlorophyll-deficient mutant), 59
 Lutescens 0026 (chlorophyll-deficient mutant), 59-61
 Lygaeids, 297
 Lyophilization procedures, 706
 Lysimeters, 168-169, 184, 186
 Lysimetric water balance studies, 189
 Lysine, 43, 655-659, 679, 700-701, 705T; and flavor, 671

M

- M 13, 83
 McEachern Jumbo (cultivar), 659T
 Macromutants, 69-70
 Macrophomina root rot, 368
 Magnesium, 147, 329; in leaf tissue, 147; research needs, 757; and yield response, 128, 147
 Magnesium content, 669T; of low quality seed, 413
 Magnesium oxide, 279
 Magnetic gates, 572
 Magnetic separation, 575T
 Main stem inflorescence, 55
 Maize, 544, 755

- Maize weevil, 521
 Makulu Red (cultivar), 100, 429, 432
 Malate synthetase, 415-416
 Malathion, 556, 744; prestorage spraying, 581; research needs, 762; resistance to, 526, 559, 562, 564-565; schedule of treatment, 557-558; spray application of, 555
 Malathion emulsifiable concentrate (EC), 554
 Malathion supplemental surface sprays, 557-558
 Maleic hydrazide (MH), 434, 436
 Male sterility, 85. *See also* Sterility
 Malimba (cultivar), 100
 Management strategies: designing of, 252-253; on individual species basis, 293. *See also* Crop Management, Production
 Mancozeb, 331, 333, 336
 Mandi (peanut synonym), 3
 Mandovy (peanut synonym), 9
 Mandubi (peanut synonym), 2
 Mandues (peanut synonym), 5, 10
 Manduiss (peanut synonym), 2
 Manduvi (peanut synonym), 16
 Manfredi Champaqui (cultivar), 336
 Manfredi 1 (cultivar), 336
 Manfredi 68 (cultivar), 336
 Manganese, 157; availability, 157; deficiency, 157; importance of, 129; tolerance of peanut to, 141; effect of toxicity, 150; yield response, 157
 Manganese content, 668, 669T; of low quality seed, 413
 Manganese ethylenebisdithiocarbamate, 329
 Manganin, 656, 657T
 Mani (peanut synonym), 1-2, 4-5, 7, 9, 16
 Mani Pintar (cultivar), 100
 Mannose, 669T
 Manobi (peanut synonym), 3, 9
 Manure, 156, 158
 Marcgrave, George, 4-5, 7-9, 11
 Marginal chlorosis, 347. *See also* Chlorosis
 Mark II, 176
 Market, peanut: description of, 747-751; domestic, 748; international, 748-750; oilstock, 590, 592, 750-751; product definition, 746-747; risks of, 751-753; sectors, 737; typical operational flow chart, 571; role of U. S. government, 737-740
 Marketing Agreement: of 1965, 378, 508, 739; of 1978, 590, 596, 598
 Marketing process: contracting, 741-743; delivery and sampling, 743-744; grading, 744; harvesting, 743; research needs, 762-764; shelling and packaging, 745; storage, 744-745
 Marketing system, 462, 571
 Mass selection, 100

- Mass spectroscopy, 673
 Material flow, 603
 Matjan (cultivar), 361
 Maturation process, 632
 Mature class, 626, 628
 Mature plant, 355-356, 363
 Mature seed: defined, 588; description of, 411; germinability, 413; grading of, 587-590
 Maturity, 114, 117; additive and nonadditive genetic effects, 80; effect on amino acid composition, 624, 657; and free arginine, 76, 630, 657; and climatic conditions, 125; dry matter accumulation and, 76; effect on fatty acid composition, 662; genotypic correlations, 76-77; and heat unit relationships, 625; heritabilities, 76-77; and mycoflora, 488-492; 497-500, 503, 505; phenotypic correlations, 76-77; research needs, 763; and seed size, 624
 Maturity methodologies, 624-625; indirect methods, 625-626; physiological maturity methods, 632-635; relative color evaluations, 626-628; research needs, 763; specific component quantitation, 630-632; weight and weight relationships, 628-630
 Maypops, 573T
 Meal moth, 521
 Meals, peanut, 702-704. *See also* Defatted peanut meal
 Mean individual kernel weight (MIKW), 628
 Meat analogs, peanuts used in, 725
 Meat loaves, peanuts used in, 725
 Meat products, peanuts used in, 725
 Meat (peanut) reclaiming equipment, 607-608
 Mechanical cultivation, 207-208, 214, 219, 243-244
 Mechanical damage, 576-579; cultivar resistance to, 132; and electronic color sorting, 618; and gravity tables, 616; and modified handling techniques, 603; from impact and pressure forces, 602; and insect infestation, 552; prevention of, 603; in receiving process, 607; during screening process, 614-616; from shelling operations, 600
 Mechanical generators, 562-563
 Mechanical shelling, 414
 Mechanical ventilation, 584-585
 Mechanics: of combining operation, 471-473; of digging operation, 460-461
 Megasporeogenesis, 68
 Meiosis: interspecific hybrids, 35-37; wild species, 34
 Meiotic chromosomes, 39-40
 Melanins, 671
 Melanosis, 338
 Melons, 504, 506
 Melting point, of peanut oil, 660
 Membrane permeability, 178, 644
 Menazon, 284, 341
 Merchant grain beetle, 534F, 536F; type of damage, 505; description of, 533, life history, 533, 535
 Mercuric chloride, 493-494, 645
 Mesocarp, 628, 634
 Mesophilic fungus, 501
 Messenger RNA, 646
 Metabolism, 415-416
 Metabolites, 486
 Metal: as foreign material, 573T, 575T; in equipment, 609
 Metalaxyl, 356
 Metham, 356, 373-374
 Metham-sodium, 362, 390-391
 Methanol, 349, 352, 627, 636, 641
 Methanolic extract, 628-630
 Methionine, 115, 655-659, 679, 700-701
 Methoxymethane, 510
 Methyl acetate, 371
 Methylamine, 510
 Methyl bromide, 559-560
 Methyl cis-3,4-dimethoxycinnamate, 331
 Methylthiopyrazine, 672
 Methyl formate, 636
 Methyl isothiocyanate, 362
 Methyl N,N'-dimethyl-N-[(methylcarbamoyloxy)-1-thioxoamimidate, 391
 Methyl 1-(butylcarbamoyl)-2-benzimidazolecarbamate, 329
 Methylpyrazine, 672, 674T, 677T
 Metolachlor, 224; nomenclature, 225T; relative toxicity of, 230T; technical description, 227; weeds controlled by, 232T
 Mice, and toxicity testing, 229, 512
 Microbial breakdown, 504
 Microbial competition, 504
 Microbial interaction, 504-505
 Microbial proliferation, 574
 Microbials, 252
 Micro-Gen® aerosol applicators, 563
 Microorganisms, 566
 Mildew, powdery, 338
 Milk curds, 719
 Milk solids-not-fat (MSNF), 723
 Miller Amendment (FDCA), 559
 Millet, 125, 512
 Milling, 590-595
 Milling losses, 575, 594
 Milling quality, 592-594; defined 115, 592; determination of 115-116; harvesting and curing operations and, 458; effects of relative humidity, 475-476; and seedcoat splitting, 115-116; effect of temperature, 475-476

- Mills:** assembly and operation, 619-620; design and operation, 602-607; edible peanut system, 602, 605F, 606F; electronic sorting, 618; gravity tables, 616-618; insect infestation and control, 541, 565; material flow and equipment arrangement, 603-607; oil processing system, 602, 604F; pre-sizing, 613-614; receiving, pre-cleaning and handling, 607-608; screening, 614-616; seed processing system, 602, 604F; seed treatment, 618-619; shelling, 608-613
- "Miltone," 721
- Mineral nutrition:** research needs, 756-757. *See also* Nutritional needs
- Minerals,** 706. *See also* specific minerals
- "Minor defects," 597T
- Minute brown scavenger beetle,** 522
- Mite predation,** 290-291
- Mites,** 503, 506, 520, 523. *See also* Spider mites
- Mitochondria,** 416
- Mitotic chromosomes,** 32F, 33F
- Mobile combine,** 474
- Modified atmosphere fumigants,** 561
- Modified atmospheres,** 602, 762
- Modified pedigree selection method,** 85, 102
- Moist heat denaturation,** 705
- Moisture.** *See* Moisture content, Soil moisture
- Moisture content,** 649; and aflatoxin invasion, 494-501; and combining time, 471; and curing, 458, 466, 470, 478; defined, 461-463; and deterioration from pests, 520; effect on drying parameter, 469; and energy consumption, 481-482; in food processing, 702, 705T; and germinability, 647-648; and milling damage, 593; and once-over method, 474; and quality losses, 474; and storage effects, 574-575, 581-582, 584, 600
- Moisture content correction factor,** 479
- Moisture equilibrium,** 495T
- Moisture meter,** 596T
- Moisture migration,** 583, 601
- Moisture retention,** 126
- Moisture transfer,** 469
- Molasses preparations,** 353
- Mold:** in grading process, 590; growth of, 554, 583; in harvesting and curing operations, 458; effect of moisture, 593; results of infestation, 595; effect of temperature, 585
- Molecular weights,** 697T, 700T
- Molybdenum,** 157-158; availability, 157; importance of, 129, 157; in nitrogen fixation process, 157
- Molybdenum (alloy),** 609
- Monardes, Nicholas,** 3, 5, 7-9
- Monkeys, and toxicity testing,** 229-230
- Monocarbonyls,** 640
- Mono-cropping,** 241
- Monogenic inheritance,** 52, 55
- Monoglycerides,** 636
- Monosaccharides,** 415, 667-668; and browning reactions, 671-672
- Monovalent salts,** 708-709
- Morning glory,** 225; proclivity, 214, 215T-216T, 218T
- Morning glory, ivyleaf,** 207, 209F
- Morning glory, smallflower,** 208T, 209F
- Morning glory, tall,** 207, 208T, 209F
- Morphactins,** 433-436
- Morphological characteristics,** 24, 220-221, 507, 755
- Morphological events,** 256
- Morphological species concept,** 24
- Morphology, effect of water stress,** 168
- Moths,** 521, 523-529
- Motor cells,** 178
- Mottled leaf mutant,** 62
- "Mouthfeel," 723
- Muck soil,** 227
- Muddy spot disease,** 332
- Multi-cropping,** 201, 241
- Multiline cultivars,** 105
- Mundubi (peanut synonym),** 7, 9
- Murky meal moth caterpillar,** 521
- Muscogee warehouse,** 578T, 584
- Mutagenic changes:** chemically induced, 70; irradiation induced, 69-70; sensitivity to, 70-71
- Mutagenicity, of herbicides,** 229
- Mutants:** chemically induced, 56; chlorophyll deficient, 59-61; in dwarfism, 55-56; radiation induced, 55; research needs, 755
- Mutation breeding,** 103
- Mutations:** from chemicals, 97; from gamma-rays, 97; as a source of genetic material, 95, 97; from X-rays, 97
- Mwitunde-3 (cultivar),** 373
- Mwitunde-7 (cultivar),** 361
- Mycoflora:** during curing in windrow and stack, 490-491; effect of inverted and random windrow drying, 491-492; before maturity and digging, 488-489; at maturity and digging, 489-490; of peanut pods and seed, 488; selective effects of the isolation medium, 494; of stored seed, 493-494
- Mycotoxins,** 486
- Mycotoxin production,** 554
- Mycotoxins,** 486, 511-513; research needs, 761; and wholesomeness, 678
- Myristic acid,** 660

N

- NAA,** 448-449
- N-butyl-N-ethyl- α,α,α , trifluoro-2,6-dinitro-*p*-toluidine,** 222
- N-methyl-N'-nitro-N-nitrosoquandine (MNNG),** 70
- N-methylpyrrole,** 672, 676T
- N,N-dimethyl-2,2-diphenylacetamide,** 221
- N³,N³-diethyl-2,4-dinitro-6-(trifluoromethyl)-1,3-benzenediamine,** 353
- N-1-naphthylphthalamic acid,** 221
- N-(trichloromethyl)thio-4-cyclohexene-1,2-dicarboximide,** 368
- Nambiquara (cultivar),** 58, 658T, 668T
- Naptalam,** 221, 224, 357; interaction with herbicides, 239-240; moisture requirement, 227; nomenclature, 225T; post-emergence treatments, 225; relative toxicity of, 230T; technical description, 227-228; weeds controlled by, 232T
- Napthaleneacetic acid (NAA),** 434, 439
- Naphtho-v-pyrones,** 513
- Natal Common (cultivar),** 388, 628
- National Peanut Council,** 601, 740
- National Peanut Research Laboratory,** 609, 613, 619
- National Seed Laboratory,** 96
- Natural enemies,** 253; of aphids, 284; of coleoptera, 303; control of insect pests, 566; and insect pest management, 251; of lepidoptera, 278-279, 296-298; of spider mites, 290-291; of thrips, 287-288; of white grubs, 307
- Natural hybrids,** 96
- Natural outcrossing,** 96
- Natural ventilation,** 584-585
- NC 2:** and effective heat units, 625; mineral element content, 669T; resistance of, 109; 352, 373; and SHMI values, 630; spider mite damage, 291
- NC 4,** 86, 421
- NC 5,** 131; amino acid composition, 659T; field losses, 470; leafspot resistant progeny, 330; pod production and growth periods, 459; and SHMI values, 630; spider mite damage, 292
- NC 6,** 131; composition of, 667T; field losses, 470; growth habit, 131; resistance of, 111, 131, 280, 288, 304; yield, 131
- NC 7:** growth habit, 131, early maturity, 114, 131; milling quality and shelf life, 131-132; resistance of, 132; yield, 131
- NC 13,** 417, 643-645
- NC 17,** 57, 291
- NC 343,** 304
- NC 3033,** 359, 387; resistance of, 105-107, 109, 358
- NC 10211,** 304
- NC 17165,** 109
- NC 17921,** 667T
- NC 17922,** 667T
- NC 17976,** 667T
- NC Ac 2575,** 343
- NC Ac 2731,** 57
- NC Ac 15753,** 304
- NC-Fla 14:** methionine content, 115; and seedcoat splitting, 116; spider mite damage, 291; evaluation of stability, 83
- NC-GP 343,** 330
- Necrosis:** from dinoseb application, 226; disease related, 330, 332, 337-338, 342, 345, 371, 375-377; and secondary invaders, 382
- Necrotic-etch leaf disease,** 71-72, 111
- Nematicides:** application of, 389-393; in disease control, 345, 361-362; in combination with fungicides, 387; effect on nitrogen fixation, 150
- Nematodes,** 187, 379F; categories of, 112-113; control of, 378-392; and crop rotation, 125-126, 241; diseases caused by, 378, 380; and fungal invasion, 487, 499, 503, 505-506; and irrigation control, 201; interaction with other pathogens, 386-387; pod breakdown and, 361-362; population dynamics, 385-386; resistance to, 43, 112-113; ring, 383; root-knot, 380-381; root-lesion, 381-383; sampling and detection, 386; sting, 383-384; and weed control, 242
- Neoxanthin,** 61
- Nervonic acid,** 661T; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T
- Net blotch,** 332-333
- Net photosynthesis (P_n):** and evaporative demand, 175; effect of increasing relative humidity, 176; and water-stress reduced chlorophyll concentration, 178; and water stress, 174-176. *See also* Photosynthesis
- Net radiant energy,** 188
- Net radiation (R_n),** 188-189; and wind turbulence, 191
- "Neutral" fractions, 483
- Neutron moderation,** 198
- Neutron-scattering technique,** 184
- New Mexico Valencia A (cultivar),** 116, 133
- New Mexico Valencia C (cultivar),** 100
- Niacin,** 670T, 684
- Nicotinic acid,** 447, 670T
- Nieremberg,** 9
- Nightshade, silverleaf,** 208T; response to herbicides, 232T; proclivity, 214, 215T, 216T, 218T
- Nighttime leaflet folding,** 178
- Nigragillin,** 513

- 9-hydroxyfluorene-(9)-carboxylic acid, . "Non-dirting" cultivation: in combination with dinoseb, 240; and control of stem rot, 552; and weed control, 217-219
- Nonfermented cheese analogs, peanuts used in, 723-724
- Nonfumigant nematicides, 391-393
- Nonpest species status, 257-258, 274
- Non-quota/additional peanuts, 738, 741
- Non-reducing sugars, 644
- Nontarget effects, 390-393
- Nontarget pest resurgence, 279-280
- Noogoora bur, 213
- Norea, 241
- Normal seedling, and germination quality, 418
- Northern root-knot nematode, 380
- North-south rows, 192-193. *See also* Row spacing
- Nubbin circuit, 607. *See also* "Raisins"
- Nuclear factors, 85
- Nuclear genes: manipulation of, 87; in chemically induced mutations, 70; interaction with plasmon, 97
- Nuclear mutations, 103
- Nuclear polyhedrosis virus, 279, 566
- Nucleic acid, 414, 649
- Nutgrasses, 573T, 575T
- Nutgrass tubers, 603, 618
- Nutritional needs, 139 et passim.; supplied through irrigation systems, 201; research needs, 755; effect on seed quality, 412-413
- Nutritional quality, of peanuts: carbohydrates, 684; minerals, 684; oil, 682-684; protein, 679-682; vitamins, 684
- Nutsedge, purple, 208T; growth factors, 222; response to herbicides, 232T; rate of occurrence, 217
- Nutsedge, yellow, 208T; 209F; growth factors, 222; response to herbicides, 232T; rate of occurrence, 217; and yield reductions, 213
- Nutsedges, perennial: and herbicide control, 222-223; problem and control of, 244; proclivity, 215T, 216T, 217, 218T
- Nyctinasty, 178
- Nylon-saran-EVA resin bags, 649
- O
- Oats, 359, 504, 506, 512
- Occasional pest status, 257, 274. *See also* Pest status
- Octanal, 636-637, 673
- Octane, 636
- o-ethyl S,S-dipropyl phosphorodithionate, 353, 391
- 9-hydroxyfluorene-(9)-carboxylic acid, 433
- 9-square bar cylinder, 609
- Nitralin, 222-223
- Nitrate reductase activity, 444
- Nitrate salt, 237
- Nitrogen: effect on *A. flavus*, 329, 496, 502; cultivar differences, 148, 151; as fertilizer, 128, 148-149, 757; effect on herbicide activity, 237; in modified atmosphere fumigation, 561; and photosynthetic rate, 151; and physiological development, 150-151; effect on quality, 412; research needs, 762; inoculation with *Rhizobium*, 149-150; relation to sulfur levels, 154; yield response, 148
- Nitrogenase activity, 445-446
- Nitrogen content, 444
- Nitrogen fixation: effect of climatic factors, 150, 445-446; role of cobalt, 150, 155; combining ability for, 80; effect of fertilization, 128; and growth regulators, 756; effect of herbicides and nematicides, 150; effect of peanut host on efficiency, 149-150; role of molybdenum, 157; research needs, 755-756; 760; by rhizobia, 104, 141, 148, 757; rhizobium strains, 443-444; effect of soil acidity, 150; strain evaluations, 444-445; response to water deficit, 180. *See also* Rhizobia
- Nitrogen solubility, 710, 723
- Nitrogen solubility index, 704, 725
- Nitrogen solubility profiles, 707-710, 711F, 712
- Nitrous acid (NA), 70
- Node development, 168
- Nodulation, 180; effect of cobalt treatments, 150; effect of herbicides and nematicides, 150; plant character, 56-57; caused by rhizobia, 141, 419
- Nodulation scores, 444
- Nodules: development, 444; leghemoglobin, 444; nitrogenase, 444; number, 180, 445; *Rhizobium* vs. nematode galls, 380-381; weights, 180, 445
- Noise, in milling process, 602
- Nomina nuda*, 22
- Nonadditive gene action, 78
- Nonadditive genetic effects, 80, 114
- Nonadditive genetic variance, 79
- Nonanal, 636-637, 673
- Nonarachin, 655, 701; composition, 656, 697-698. *See also* Conarachin
- Nonchemical pesticide control, 387-388, 759
- Nondestructive maturity method, 633-634

- Off-flavor, 477
- Oil, peanut, 571, 750; aflatoxin removal, 508, 510; color, 626-627, 634; composition of, 660-666; content, 114, 634, 655, 660; in curing process, 458, 636; extraction of, 14, 656, 703; in Europe, 14, 16-17, 750; and herbicides, 240; inheritance studies, 75-77, 79-80, 114-115; and irrigation, 174; nutritional quality, 682-684; price levels, 750-751; research needs, 755; stability of, 84, 664-665; in United States, 17, 689, 751
- Oil absorption, 721
- Oil-base spray: formulation, 558; as surface spray, 556; toxicity, 557
- Oil mobilization, 704
- Oil processing: operational flow chart, 590F; primary use, 590, 592
- Oilseed flours: baking properties of, 726; bread-baking properties, 727; in bread formulas, 726
- Oilstock, 750-751; Segregation III peanuts, 508, 590, 750-751; Segregation II peanuts, 750-751; values, 750
- Oilstock market, 592
- Oil synthesis, 257
- Okra, 373
- Okra mosaic virus, 347
- Oleate-linoleate ratios, 663T, 664T, 665
- Oleic acid, 660-664, 682; mature vs. immature seed, 662; and oil stability 662, 664; roasted vs. unroasted peanuts, 666T triglycerides and phosphatides, 665T
- Oligosaccharides, 415, 667-668
- Olive oil, 683
- Olsen extractant: calcareous vs. noncalcareous soil, 152-153; composition, 152; critical level, 152
- o-methylsterigmatocystin, 512
- Once-over harvesting procedure, 474
- 1,1-dimethyl-4,4-bipyridinium ion, 353
- 1,1-dimethyl-3-(d,d,d-trifluoromethyl)urea, 353
- 1,3-D, 389
- 1,3-dichloropropenes (DD;1,3-D), 389
- 1,2-dibromo-3-chloropropane, 240, 389
- 1,2-dibromomethane (ethylene dibromide; EDB), 389
- o,o-diethyl o-[p-(methylsulfinyl)phenyl]-phosphorothionate, 353, 391
- Optical density, 628
- Optimum day/night temperatures, 430
- Optimum harvest time, 632
- Optimum moisture, 136
- Oral LD₅₀, 229T, 230T
- Organic acids, 636
- Organic fertilizers, peanuts used as, 510
- Organic matter content, 367, 551; and crown rot, 367; effect on herbicides, 238; importance of, 242; effect on mycoflora, 492, 506; and soil type, 238
- Organic mercurial fungicides, 367-368
- Organochlorine compounds, 291, 303
- Organoleptic quality, 722-724, 730-731; evaluation of, 640
- Organophosphorus (OP) insecticide, 291
- Origins, of cultivated plant, 1-2, 11-13, 95; center of, 51F, 52; from diploid ancestors, 32
- Osmophilic fungi, 492, 494
- "Other kernels," 173-174, 588
- Overflood irrigation, 292
- Overhead belt conveyor, 579, 581
- Overmature plants, 369
- Overmature seed, and aflatoxins, 499-500
- Overo, 336
- Overspace mechanical ventilation system, 583-584
- Ovule culture, 447
- Oxidative stability, 636
- Oviedo y Valdés, Captain Gonzalo Fernandez de, 1-2
- Oxadiazon: nomenclature, 225T; preemergence treatments, 223; relative toxicity of, 230T; effect on stem rot, 240-241, 353; technical description, 228; weeds controlled by, 232T
- Oxalate extractant: composition, 152; critical level, 152-153
- Oxalic acid, 349, 356
- Oxamyl, 391
- Oxazoles, 673-674, 678
- Oxazolines, 673-674, 678
- Oxidative rancidity, 114
- Oxygen (O₂) concentrations: and germination, 414; effect on apparent photosynthesis, 424; and growth of *A. flavus*, 496, 501-503
- Oxygen solubility, 502-503
- P
- Packaging of peanuts, 745-746
- Pairing relationships, 39-40
- Palmitate, 665
- Palmitic acid, 660, 661T-663T; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T
- Palmitic acid, 661T, 665T
- Panicum, fall, 208T, 212F; response to herbicides, 232T; proclivity, 214, 215T, 216T, 218T; and yield reductions, 213-214
- Panicum, Texas, 208T, 209F; response to herbicides, 222, 228, 232T; proclivity, 214, 215T, 216T, 218T
- Pantothenic acid, 670T
- Papain, 710, 717, 719

- Paradichlorobenzene, 38, 648
 Paraquat, 353
 Parasites, 522, 566, 762; control of *A. gemmatilis* larvae, 279; of *E. lignosellus*, 296, 297T; of lepidopteran larvae, 278; of *S. rolfii*, 353; of thrips, 287-288; of white grubs, 307; as suppressive agent, 251-253; and survival of *R. solani*, 363
 Parasitic fungi, 492. *See also* Fungi
 Parkinson, 4-5, 9
 Partitioning: to fruits, 168-169, 758; of photosynthate, 432
 "Partitioning coefficient," 432
 "Paspalum staggers," 512
 Pathogenic fungi, 487. *See also* Fungi
 Pathogens, 566, 705T; in integrated pest management, 251; in pest control, 253; as vectors, 287
 Patulin, 513
 PCNB, 240; control of charcoal rot, 369; control of *S. rolfii*, 352-353, 392-393
 p-coumaric acid, 643
 p-hydroxybenzoic acid, 643
 PCP, 219, 221
 Pea, 340
 Pea enation virus, 341
 Peak evapotranspiration, 185. *See also* Evapotranspiration, Potential evapotranspiration
 Peak leaf area index, 185. *See also* Leaf area index
 Peak ratio analysis, 641
 Peak water use periods, 137. *See also* Water use efficiency
 Peanut Administrative Committee (PAC), 508, 739-740
 Peanut butter: acceptance of, 738-739, 749-750; Canadian manufacturers, 747; choline content, 670; domestic food use, 655, 689; flavor, 671; effect of malathion treatments, 554; first patent, 17; use of split seed, 745-746; viscosities, 671
 Peanut chlorotic leaf streak, 347-348; causal agent, 348
 Peanut chlorotic leaf streak virus (PCLSV), 347-348
 Peanut clump, 344; causal agent, 344-345; control of, 345; epidemiology, 345; symptoms and host range, 344
 Peanut clump virus (PCV), 344-345
 Peanut combine: and broadleaf weed damage, 244; cut-away view, 472; development of, 17-18; effect of growth regulators on performance, 136
 Peanut dispenser, penny machine, 17
 Peanut green mosaic, 346; causal agent, 346
 Peanut green mosaic virus (PGMV), 282, 346
 Peanut Insect Defoliation Model, 277-278
 Peanut Loan Schedule, 588, 589T
 Peanut Marketing Agreement (1978), 590, 596, 598
 Peanut milk systems, 721-722
 Peanut mottle, 339; causal agent, 339; control of, 340; epidemiology, 339-340; symptoms of, 339
 Peanut mottle virus (PMV), 282T
 Peanut picker, patented, 17
 Peanut polisher, 15
 Peanuts: conceptualization of system, 252-253; dispersion of, 13-15, 754; domestication of, 15; early history, 1-11; enemies of, 253; ethnological comparisons, 15; history, 3, 13-14, 754; Indian vernacular names, 9, 16; leaf anatomy, 419-420; linguistic affinities, 16; origin of, 1-2, 11-13, 32, 51-52, 95; uses of, 13, 16-18, 510, 590, 592, 655, 739, 745-747, 750. *See also* Cultivars, Cultivated peanut
 Peanut sheller, 14, 115
 Peanut stunt, 343; causal agent, 343-344; control of, 344; epidemiology, 344; resistance to, 43; symptoms and host range, 343
 Peanut stunt virus (PSV), 282T, 343-344; vectors of, 287
 Peanut yellow mottle, 347; causal agent, 347
 Peanut yellow mottle virus (PeYMV), 347
 Pearl (cultivar), 641
 Peat moss, 238
 Pebulate, 222-223
 Pectin methyl esterase, 647
 Pedigree selection method, 85
 Pedigree natural crossing, 96
 Pegging zone: importance of calcium, 128-129, 139, 141; effect of temperature, 443
 Pegging zone moisture content, 171
 Peg rot, 128
 Pegs: behavior of, 24, 135; and diseases, 363, 365; elongation, 183; effect of growth regulators, 434-435; moisture requirements, 181-183; production of, 431; effect of relative humidity on growth rate, 441; effects of soil water deficit, 169-171; and optimum water management, 201
 Pelargonidin, 68
 Pennsylvania smartweed. *See* Smartweed, Pennsylvania
 Pentachloronitrobenzene (PCNB), 240, 352-353, 369, 392-393
 Pentachlorophenol (PCP), 219, 221
 Pentanal, 636, 673
 Pentane, 636-639, 641
 Pentose-phosphate pathway, 415
 Peonidin, 68
 Pepper spot, 334; causal organism 334; control of, 335; disease cycle and epidemiology, 334-335; symptoms of, 334
 Pepsin, 680, 710, 716-717, 720

- Peptides, 656
 Percolation, 184, 186, 192, 194
 Perennial growth habit, genetics of, 55
 Perennial weeds, 758. *See also* names of individual weeds, Weeds
 Perforated sheet metal type commercial sheller, 609, 612T
 Pericarp, 64, 72; accumulation of starch, 415; colonization by *A. flavus*, 489; changes in, 633-634; color of and maturity, 626; correlation coefficients of yield, 105
 Peroxide content, 636
 Peroxides, 647
 Peruvian cultivar type, 12-14, 51
 Pestalotiopsis leafspot, 338-339
 Pest densities, 257, 259
 Pest dynamics 271
 Pesticide residues, 562
 Pesticides: and cultivar interaction, 239-241; and genotype interactions, 113; in insect control, 520 et passim.; integrated pest management, 250, 253; and irrigation practices, 164, 201; and wholesomeness, 678. *See also* Insecticides
 Pests: control of, 552-566; crop-pest relationships, 253-257; damage caused by, 254-255; hierarchies of, 252-253; research needs, 756, 758-759, 762; resistance to, 43, 71-72, 106; effect on storage, 552-553; tolerance of plant to, 255. *See also* Insects
 Pest status, 271; factors affecting, 258; types of, 257
 pH, and growth of pathogens, 328, 355, 361, 372
 pH, of seed, 694, 708, 713-716, 718-721
 pH, of soil, 157, 231, 237-238, 242
 Phenamiphos, 391
 Phenolic acids, 363, 507, 643, 650
 Phenological periods, 280
 Phenological stages, 181
 Phenols, 674T
 Phenotypic coefficient of variation, 74-75
 Phenotypic correlations: broad-sense heritability, 75-76; and early maturity, 114; for fruit and seed volume, 105; vs. genotypic correlations, 75, 85; and heritability, 75; for yield, fruit size and maturity, 76-77, 104
 Phenotypic stability, 83
 Phenox herbicides, 219
 Phenylacetaldehyde, 673
 Phenylalanine, 655-659, 701; and flavor, 671
 Pheromones, 762
 Philippine White (cultivar), 58, 72
 Phlobophene, 68
 Phloem sap, 179
 Phloem tissue, and calcium translocation, 142, 146, 172
 Phomopsis leaf and stem diseases, 337-338
 Phorate, 240
 Phosphatase, 636
 Phosphatides, 665T, 666
 Phosphatidylethanolamines, 665
 Phosphatidyl serines, 665
 Phosphine, 559-560
 Phosphorus, 151-153; in combination with nitrogen 148; deficiency of, 151; effect on evapotranspiration and water use efficiency, 193; types of extractants, 152; and late leafspot, 329; effect on quality, 412; research needs, 757; and high yields, 128
 Phosphorus content, 669T; of leaf tissue, 153; of low quality seed, 413
 Photoelectric color sorting, 509
 Photoperiod, 440-441, 760; and flowering, 440; effect on legume nodulation, 446; temperature interaction, 440
 Photorespiration, 424
 Photosynthate, 170-171, 183, 432, 760; production of, 256
 Photosynthesis, 43; and diffusive resistance, 174-178; and evaporative demand, 190; and growth regulators, 756; characteristics of leaf, 419-421; effect of leaf and plant age, 424-425; of plant canopies, 425-427; research needs, 755; response to temperature, 421F, 430; effect of water deficit, 183. *See also* Chlorophyll
 Photosynthetic capacity, 421
 Photosynthetic photon flux density (PPFD), 177F, 189
 Photosynthetic rates: and leaf-nitrogen concentration, 151; effect of transpiration, 191; and yield stability, 104
 Phyllosticta leafspot, 338
 Physicochemical properties, of peanut proteins, 689-701, 763
 Physiological activity, and drought stress, 499-500
 Physiological deterioration, 411
 Physiological events, 256
 Physiological maturity, and digging time, 459
 Physiological Maturity Index (PMI), 628, 633
 Physiological maturity methods, 632-635
 Physiological relationships, 59, 61-62
 Physiological traits, 80, 755, 762
 Physiology, and plant nitrogen, 150-151
 Phytoglycolipids, 666

- Phytohormones, 438
 Phytophagous arthropods: classes of, 259;
 foliage inhabiting, 271-292; in manage-
 ment strategies, 258; soil-inhabiting,
 292-307; table of, 260-270
 Phytotoxic interactions, 237 et passim.
 Phytotoxin, 371
 Phytotron unit, 102
 Picking efficiency, 472
 Pickup operation, 472
 Picloram, 434
 Pic + T, 434, 437
 Pigment production, inheritance of, 66-68
 Pigmy seed, 64
 Pigweed, 208T, 212F; response to her-
 bicides, 225, 232T; proclivity, 214,
 215T, 216T, 218T
 Pigweed, redroot, 207, 215T
 Pigweed, smooth, 213
 Pindalls (peanut synonym), 10
 Pineapple beetle, 522
 Piperonyl butoxide, 554, 556, 558, 563
 Piso, Willem, 7
 Pistache (peanut synonym), 9, 11
 Piston-type pump, 555, 557
 Planaltine ellipse, 25, 27F
 Plant characters: branching patterns, 50,
 53, 55; cytoplasmic inheritance, 85-87;
 dwarfism, 55-56; genetic linkages, 72;
 growth habit, 52-54; and irrigation prac-
 tices, 193; main stem inflorescence, 55;
 and maturity, 55; nodulation, 56-57
 Plant Growth Regulator Working Group,
 437
 Planting: recommended types of beds, 133;
 dates of, 133; effect of density on evapo-
 transpiration, 192-193; drill spacing 134;
 row spacing, 134
 Planting patterns, 134; and disease control,
 329; and weed control, 242
 Planting time, 126; and disease control,
 301, 329, 341, 343, 374; and pest con-
 trol, 251, 285
 Plant introduction, 95-96, 99-100, 388,
 754-755
 Plant Introduction (PI) numbers, 96, 98.
See also individual numbers
 PI 109839, 107-108
 PI 123643, 86
 PI 159747, 340
 PI 161317, 132
 PI 196613, 281
 PI 203395 and PI 203397, 107
 PI 221068, 304
 PI 259747, 57
 PI 261893-2-1-1-B, 107; -1-3-B, 107; -2-1-
 4-1-2B, 107
 PI 2619061-1-1-1-2-B-B, 107
 PI 261942 and PI 261943, 72
 PI 261945 and PI 261946, 340
 PI 262286, 292
 PI 262827, 292
 PI 262840, 292
 PI 262848, 343
 PI 269685, 107
 PI 280688, 112, 288
 PI 290599, 288
 PI 290606, 112-113, 388
 PI 295233, 112-113, 388
 PI 337394F, 110
 PI 337409, 110
 PI 338296, 292
 PI 338317, 292
 PI 341884, 377
 PI 341885, 110
 PI 341886, 377
 PI 341995, 377
 PI 343360, 110
 PI 365553, 110, 507
 PI 371521, 356
 PI 372263, 106
 PI 372303, 106
 Plant turgor pressure, 183
 Plasma lipids, 683
 Plasmogenes, 70, 87
 Plasmon: constitution, 86; differentiation,
 40; genetic variability in, 97; inheritance
 of growth habit, 53; mutations, 103
 Plastochron, 168
 Plenum, 477, 480
 Ploidy alteration, 97-98
 Ploidy level manipulation, 44-45
 Plukener, Leonard, 4, 10-11
 Plumier, Charles, 4-5, 10-11, 14
 Plumule damage, 413. *See also* Dark
 plumule
 Pneumatic conveying, 472
 Pneumatic loaders, 579
 Pneumatic samplers, 586-587, 744
 Pod, 487, 595; cytoplasmic inheritance, 86;
 inheritance of constriction, 64; rates of dry
 matter accumulation, 429; early reports,
 5-6; initiation of, 254, 432; maturity and
 digging losses, 460; production, 347,
 459; pubescence, 64-65; retention, 435;
 reticulation, 65; classification of shape,
 594F; effect of temperature on growth,
 430-431; types of, 65; effect of water de-
 ficit, 169-170; water potential, 179;
 weight, 75-76, 429
 Pod breakdown, 105; and aflatoxin con-
 tamination, 488-492; caused by disease,
 359-361, 363, 365-366; and soil calcium
 level, 145-146

- Pod characters, 64-65, 130
 Pod development: and AMI, 630; and
 fungi, 486, 489-490; effects of light on,
 439; moisture requirements, 182-183;
 and soil moisture content, 170-171
 Pod fill: calcium requirement, 756; duration
 of, 432, 756; and irrigation, 133
 Pod formation: and drought sensitivity,
 182-183; effect of growth regulators, 435-
 437; and optimum water management,
 201; effect of temperature on, 430-431
 Pod Maturity Profile, 628, 634-635
 Pod number, 65, 429, 431; correlation coef-
 ficients, 75-76; effect of cultivation, 135;
 dominance effects, 75; general combining
 ability, 79; heterosis, 78; inheritability,
 75; nonadditive genetic effects, 81; and
 soil water deficit, 170; effect of tempera-
 ture, 441
 Pod rot: caused by diseases, 349, 351F, 354,
 359-361, 363, 365, 372; due to fertiliza-
 tion, 128
 Pod set, and Shellout Method, 626
 Pod size, 131-132; combining ability, 80;
 heritability, 77; heterosis, 78; inheritance
 of, 64
 Pod surface fungi, 491
 Pod yield: combining ability, 79-80; domi-
 nance effects, 75; in epistasis, 81; genetic
 advance, 77; heritability of, 73-77; in
 heterosis, 78; nonadditive genetic effects,
 81; and thrips infestation, 287
 Poikilotherms, 271
 Polarized laser beams, 509
 Polenske value, 660
 Pollution, in milling process, 602
 Polyacrylamide disc gel electrophoresis,
 656, 690, 692; patterns of, 694F, 699F
 Polyethylene liners, 601
 Polymers, high density, 609
 Polypeptide chains, 695-696
 Polypeptides, 656-657, 698
 Polyploidy, 30, 39-40, 45, 98
 Polysaccharides, 706
 Polyunsaturated fatty acids, 682
 Polyunsaturated oils, 682
 Polyvinyl chloride strips, 554, 563
 Poona Local (cultivar), 57
 Poona White (cultivar), 57
 Population density, 257, 289, 552
 "Pops," 630; calcium deficiency and require-
 ments, 128, 142, 144; as foreign mate-
 rial, 573T, 575T; from water stress in
 fruiting zone, 172
 Population dynamics, 385-386
 Porometer, 176
 Postemergence herbicide applications,
 135
 Postharvest chemical treatments, and dor-
 mancy 644
 Postharvest insect control, 762
 Postharvest physiology, 624, 635; curing,
 635-636; dormancy, 641-647; flavor,
 636-641; research needs, 762-763; stor-
 age, 647-650
 Potassium, 146-147; in crop rotation, 146-
 147; deficiency of, 147; in fruiting zone,
 145, 412; effects on herbicides, 237; and
 late leafspot, 329; and leaf display, 178;
 research needs, 757; effect on yields, 128
 Potassium azide, 352
 Potassium content, 669T; in low quality
 seed, 413
 Potato leafhopper, 112-113, 131
 Potato virus Y group, 339, 346
 Potatoes, 504, 506, 512
 Potential evapotranspiration (E_0), 183,
 185
 "Potyvirus group," 346
 Poultry, 512
 Powder fertilizer, 128, 145
 Powdery mildew, 338
 PR 64B (Malawi), 87
 Precision planting, 126
 Precleaner, 572, 607; operational flow
 chart, 576F
 Precleaning, 607-608; equipment, 572,
 580T
 Predaceous phytoseiid mites, 290
 Predators: of insects, 566; in integrated pest
 management, 251, 253; of pests, 287-
 288, 297; research needs, 759, 762
 Predigging operations, 460
 Preemergence seedling rot, 375
 Preemergence treatments, 220
 Preharvest conditions, effect on germina-
 bility, 412-413
 Preharvest insect management, 758-759.
See Integrated pest management
 Preharvest insects: crop-pest relationships,
 253-257; pest management, 250-253;
 pest status, 257-259
 Preharvest pod rot, 362
 Preparation of land. *See* Land preparation
 Preplant incorporated (PPI) herbicides,
 127, 135
 Prepress solvent extraction, 702
 Prepressing, 702-703
 "Prescription approach," 244
 Presizer, 586-587
 Presizing, 607, 613-614
 Pressure bomb technique, 179
 Pressure drops, 480
 Prevention, of insect pests, 251-252
 Price support system, 586; influence of,
 571; and irrigation, 164; future role of,
 757, 764; of U. S. government, 738-740,
 749
 Primary dormancy, 642

- Processing plant: and insect infestation, 535, 537, 541, 543, 549, 552; insecticidal aerosol space treatments, 561-564; effect of temperature extremes on insects, 565
- Processing system, 571
- Processors, peanut: cost reduction, 746-747; markets and market risks, 747, 752; needs of, 95, 117, 746
- Procymidone, 356
- Producers, peanut, 764; capital requirements, 458; and insect infestation, 552; needs and desires, 95, 117; profit maximization and digging time, 459; use guidelines, 586
- Product definition, 746-747
- Production: fixed area for, 17, 164, 737, 741, 764; costs of, 103, 641, 756, 764; of edible peanut products, 701-707; and growth regulators, 135; and herbicides, 244; management, 412; price support systems, 571, 738, 740-741, 749, 757, 764; water requirements, 186; in the United States, 123-124, 737-740, 750-751, 763-764; effect of weeds, 134, 214
- Product utilization, and storage, 647
- Profile available water, 192
- Profile water depletion, 200
- Profluralin, 236
- Progeny management, hybrid, 101
- Proline, 657T-659T, 701; in arachin, 655; and drought stress, 180-181
- Prometryn, 222
- Pronto (cultivar), 133
- Propeller-type fans, 585
- Prophylaxis, 280
- Propyl gallate, 638
- Protease, and curing, 636
- Protectants, 555-556
- Protein, 29, 43, 415, 655; and aflatoxin contamination, 511; amino acid levels, 115, 656, 657T; components, 655; in dairy applications, 722; emulsion formation, 713-717; extraction and fractionation of, 689-701; flow properties, 713; foam formation, 717-719; gelation, 719-720; isolation of, 706; nutritional quality, 679-682; oil absorption properties, 721; range of, 655-657; recovery, 693-694; solubility, 647, 707-712, 719, 723; sources of, 708, 713; synthesis, 415-416; water sorption, 721
- Protein concentration, 708, 713
- Protein content: combining ability, 80; of peanut flour, 705T; of peanut fruit, 429-430; heritability of, 76-77; and irrigation 174, phenotypic correlations, 77; research needs, 755, 763; effect of sulfur fertilization, 154
- Protein efficiency ratio (PER), 679, 723
- Protein isolates, 723, 727
- Proteolysis, 693
- Protocatechuic acid, 643
- Protoplast fusion, 45
- Protozoans, and insect control, 566
- Provitamin A, 670
- Prudente INTA (cultivar), 336
- Pseudoplastic flow behavior, 722
- Pseudoscorpion, 523
- Psychrometric chart, 476, 477F
- Pure line selection, 99-100, 102
- Purple crescent, 57
- Purple nutsedge, *See* nutsedge, purple
- Purslane, common, 208T; response to herbicides, 232T; proclivity, 214, 215T
- Pusley, Florida, 208T, 210F; control of, 244; response to herbicides, 221, 225, 232T; proclivity, 214, 215T, 216T, 218T; and yield reductions, 213
- Pyrazines, 672-675, 677T-678T
- Pyrethrins, 554, 556, 558, 563
- Pyriclor, 222
- Pyridines, 674T, 677T
- Pyridoxine, 447, 670T
- Pyrroles, 672-673, 675T-677T
- Pythium diseases, 359-360; causal organism, 360; control of, 361-362; crop rotation and, 126; disease cycle and epidemiology, 360-361; symptoms of, 360
- Pythium pod rot, 109-110, 351F, 755; control of, 361-362; and soil calcium levels, 145-146; symptoms of, 360

Q

- Qualitative characteristics: of flower, 57-63; of induced mutagenic changes, 69-71; of linkage, 72; of pest resistance, 71-72; of plant, 52-57; of pod and seed, 64-69
- Quality: effect of curing, 458; effect of fumigation, 560-561; and insect contamination, 520, 523, 529-530; and irrigation, 136; maturity methodology and, 624; milling, 115-116; of oil content, 114-115; of protein content, 115; and rainfall, 125; research needs, 760, 762-763; effect of soil minerals, 128-129, 154; storage effects, 574; 640, 647-649; role of USDA, 739; effect of water deficit, 173-174
- Quality Committee of APREA, 678
- Quality evaluations, 482-483
- Quantitative characteristics: effect on breeding procedures, 84-85; cultivar stability, 83-84; gene action, 73, 77-82; genotype x environment interactions, 82-83; germplasm pool diversity, 73-75; heritability, 73, 75-77; yield, 72
- Quiescence, 642
- Quinones, 529, 533

- Quota peanuts: and *A. flavus*, 744; concept of, 738; defined, 741; domestic market, 748; international market, 748; price support system, 744

R

- Rabbits, in toxicity testing, 229-230, 683
- Radiant energy supply, 188, 190
- Radiation, 189; intensity, 189
- Radiation-induced mutants, 55, 59, 62, 69-70
- Radioactive lipids, 636
- Raffinose, 415, 666-668, 704
- Ragweed, 208T; response to herbicides, 232T; proclivity, 214, 215T, 216T, 218T
- Rainfall: climatic requirements, 125; and cumulative pan evaporation, 200; distribution of, 125; effects of, 469-470; effects on herbicides, 233-236; importance during pod fill, 133; and irrigation scheduling, 195; and peanut production, 186; and pest status, 258; and water account, 200; growth of web blotch, 333; and yield reduction, 183
- "Raisins," 572, 573T, 574T, 594, 630
- Rancidity, 574, 590
- Random windrow drying, 468, 491-492, 499
- Rats, and toxicity testing, 229, 512
- Rapeseed oil, 660
- Ray, John, 4, 9, 11
- Receiving, 607-608
- Recirculating sprayer, 224
- Recognition of pests, 520-521, 523 et passim.
- Recombination, 100-101
- Recommended dietary allowances (RDA), 681
- Red flour beetle, 531F, 532F; type of damage, 530, 533; description, 530; life history, 530; resistance to malathion, 565
- Redhorned fungus beetle, 522
- Rednecked peanut worm, 391
- Red pigment production, 57
- Reducing sugars, 644, 649, 704
- Reductive pentose phosphate (C_3) pathway, 420
- Reel screeners, 614-616, 616T
- Reflectance spectrophotometry, 671
- Refractive index, 660T
- Refrigeration, 648; in insect control, 565; in storage, 601
- Rehovot 121 (cultivar), 109
- Reichert-Meissl value, 660
- Relative color evaluation, and crop maturity, 625-628
- Relative humidity, 475; and aflatoxin invasion, 378, 494-501, 504; effect on artificial drying, 475-477; and bin drying, 478-479; equilibrium moisture content, 463, 465, 475; effect on evapotranspiration, 190; effect on growth, 441-443; optimum for insect infestation, 258, 526, 535, 537, 543-544, 547, 551-552; effect on lipids and nucleic acids, 414; effects on milling quality, 476; and storage effects on dormancy, 646, 648; effect on stored seed, 414, 582; temperature effects on, 468; and water sorption characteristics, 720; and windrow drying, 467. *See also* Humidity, Moisture, Soil moisture
- Relative leaf water content (RWC), 174, 179
- Relative pod maturity, 633
- Repellants, 521
- Reproductive growth: flower formation, 429; effect of growth regulators, 434-437; rate of pod dry matter accumulation, 429; pod growth, 429; pod number and weight, 429; research needs, 760; effect of temperature, 443; and water stress, 168
- Reproductive stages, 181
- Reproductive tissue, culture of, 448-449
- Required drying time, 478
- Research Institute for Oils and Oilseeds, 180
- Research needs, 754-764
- Residual fertility, 125, 128, 135
- Residual insecticides, 554
- Residual spray, 554
- Residues. *See* Crop residues
- Residue tolerances, 559, 562
- Resistance, to air flow, 479
- Resistance meters, 137
- Resistance traits: to foliar diseases, 330, 332, 334-336, 340-341, 343, 345; to nematodes, 387-388; research needs, 755; to soilborne diseases, 352, 356, 358-359, 361, 364, 366-367, 369, 371-374, 377-378
- Resistant cultivars/germplasm: to aphids, 284-285; to coleoptera, 304-305; to foliage consumers, 280; integrated pest management and, 251, 253; to lepidoptera, 280-281, 300; to spider mites, 291-292; and control of stem rot, 352; to thrips, 288
- Respiration, 635-636, 644
- Respiratory activity, 416
- Respiratory quotient (RQ), 635

- Retention indices, 673
 Reverse osmosis, 706
 RG-1, 103, 111
 Rhizobia: and acid-soil infertility, 141; and need for cobalt, 155; sensitivity to heat and drought, 180, 760; and nitrogen fixation, 104, 128, 148, 180, 757
 Rhizobium strain, 443-444, 446; evaluations of, 444-445
 Rhizoctonia diseases, 362; causal organism, 363; control of, 364; disease cycle and epidemiology, 363-364; symptoms of, 362-363
 Rhizoctonia foliage blight, 337, 759
 Rhizomes, as foreign material, 573T, 575T
 Rhizopus diseases, 375; causal organism, 375; control of, 375; disease cycle and epidemiology, 375; symptoms of, 375
 Rhizosphere, 367, 488
 Riboflavin, 330, 670T
 Ribosome system, 416
 Rice, 176, 512, 544, 549
 Rice moth, 528F; type of damage, 529; description of, 528; life history, 528-529
 Rice weevil, 521
 Ridge cultivation, 217
 Ridomil, 356
 Ringlegged earwig, 522
 Ring nematodes, 379F, 383, 389-393
 Ringspot disease, 341-342
 "Ripple-type" grate, 613
 RMP 12, 341
 RMP 91, 341
 RNA: content, 650; synthesis level, 416. *See also* Messenger RNA
 Roasted peanuts, 748-749
 Roasters, needs of, 746
 Roasting, 679-680; and aflatoxin reduction, 510; effect on sugar content, 669T
 Robur 33-1, 114, 343
 Rocks/stones, 573T, 575T, 618
 Roll feeder, 572
 Ronstar, 225T
 Rootbed, 126
 Root-cell membrane resistance, 146
 Rooting zone, 195; and available free water, 170-171, 193
 Root-knot nematode, 379F, 380; and crop rotation, 126, 388; infection and spread, 381; nematicide control, 389-393; resistance to, 113, 388; symptoms of, 380-381. *See also* Nematodes
 Root-lesion nematodes, 381; and crop rotation, 126; infection and spread, 382-383; resistance to, 112-113; symptoms of, 381-382. *See also* Nematodes
 Root rot: causes, 354, 359-363, 365; resistance to, 361
 Root rot disease, 368
 Roots: depth progression, 165; extension of and water extraction, 181, 758; growth of, 429; length of, 165, 166F; tissue culture, 449; types of, 24; and water uptake, 165-167; weight of, 165
 Rosette virus, 340; causal agents, 341; control of, 341; epidemiology, 341; resistance to, 43, 71, 103, 110-111, 341; symptoms of, 340-341
 Rotary hoe, 219
 Rotation time, 201. *See also* Crop rotation
 Rough testa, inheritance of, 65
 Roundup, 225T
 Rovral, 356
 Rowland, Thomas B., 15
 Row spacing, 134; effect on evapotranspiration, 192-193; and integrated pest management, 251; research needs, 758; and water use efficiency, 193; and weed control, 219, 242-245
 Rugose leaf curl (RLC), 282, 347
 Runner type (cultivar), 14, 32, 53, 367, 763; calcium requirement, 129; carbonyl compounds, 640, 674T; cleaned inshell peanuts, 599; consumption patterns, 738; daily water requirements, 136; genetic studies, 101-103; geographic areas of production, 123, 132, 738; germinability and dormancy, 412-413; 644-647; gibberellin activity, 433; and grade requirements, 586, 588, 595, 597T; market varieties, 132; milling quality, 115; and planting conditions, 133-134; pod shape, 594F; resistance of, 102-103, 108, 255-256; segregation of, 576; shelling and separating, 745; stability of oil, 665; effect of temperature, 441
 Runoff, 184
 Rust, peanut, 106, 326, 330-332, 350F; causal organism, 330-331; control of, 331-332; disease cycle and epidemiology, 331; effect on evapotranspiration, 191; foliar disease complex and, 332; resistance to, 43, 71, 108; symptoms of, 330-331
 Rusty grain beetle, 521
 Rye: and crop rotation, 126, 504, 506; control of CBR, 359; and drought resistance, 113

S

- Sack drying, 483
 SADH (growth retardant), 433
 Safflower oil, 682-683
 Sakaguchi reaction, 631
 Salt concentrations, 715-716
 Salted peanuts, 748-749
 Salters, needs of, 746
 Samaru (cultivar), 190

- Samaru-38 (cultivar), 154
 Samaru 61 (cultivar), 500
 Sample trier, 596T
 Sampling methods: aflatoxin contamination, 508-509; cleaned inshell and shelled peanuts, 595-600; insect infestation, 762; for nematodes, 386. *See also* Detection
 Sand, 193, 238
 Sandbur, 208T; response to herbicides, 232T; proclivity, 214, 216T, 218T
 Sandy loam soil, 193, 196-197, 239
 Sandy soil, 213, 225-227; and pest status, 258, 423
 Sanitation, 553-554; disease control and, 285, 288, 301, 359; in edible peanut processing, 602, 607; importance of, 554; and modified atmosphere handling, 602; pest control and, 520
 Saponification value, 660
 Saprophytic fungi, 487, 492, 507
 Sap transmission, of diseases, 341, 347
 Saturated acids, 682
 Sawtoothed grain beetle, 534F, 536F; type of damage, 533; description of, 533, life history, 533
 Scab, 336; causal organism, 336; control of, 336; symptoms of, 336
 Scale sampler, 596T
 Scarabaeids, 301
 Scarabs, 306
 Schiff base cation, 672
 Schmidt, Ulrich, 2-3
 Schwarz 21 (cultivar), 21; resistance of, 109, 361, 373, 377
 Sclerotinia blight, 109, 326, 351F, 354, 759; causal organism, 354-355; chemical control of, 329-330, 356-357; and crop rotation, 126; disease cycle and epidemiology, 355-356; resistance to, 72, 131; symptoms of, 354
 Sclerotium rot, 348-349, 351F, 352-354
 Scorch, 350F. *See* Leaf scorch
 Screening, 575T, 614-616; equipment, 573
 Screen shaker, 596T
 Screw pressing, 702-703
 Seasons: effect of changes on canopy AP, 427; effect on evapotranspiration, 184-185; effect of length on transpiration, 191; trends in water use, 184, 185F, 186-187, 192
 Secondary dormancy, 642. *See also* Dormancy
 Secondary pest status, 257; as foliage consumers, 274. *See also* Pest status
 Second-order interactions, importance of, 83
 Sedges, 208T; response to herbicides, 227; problem of, 217. *See also* Nutsedges
 Sedimentation coefficients, 690
 Seed, 5, 6F, 487, 591F, 618; amino acids, 657; and AMI values, 631; calcium level, 174; effect of cultural practices, 411-412; damage, 471, 475, 493, 503-504; decay of, 375; disease transmission, 342, 346; dormancy, 65, 102, 416-417, 635, 641-647; drought sensitivity, 183; energy value, 679; effect of growth regulators, 435-437; and herbicide injuries, 241-242; insect infestations, 287, 520; maturity of, 254, 256, 624; moisture, 413-414, 493, 648, 703; oil content, 655, 660, 679; and physiological maturity, 632; planting conditions, 134, 242-243; processing of, 704, 745; protein content, 29, 43, 76-77, 80, 154, 174, 655 et passim. 679; quality of, 241-242, 412, 508-509; shape, 65; size, 65, 72, 78; storage and storage effects, 414, 647; uses of, 590, 592; weight, 75; yield, 75, 77, 79
 Seedbed preparation, 126, 133, 206, 217-219
 Seed characters: arginine content, 69; dormancy 65; importance of, 104-105; shape, 65; size, 65; testa color, 65-68; inner testa color, 69; rough testa, 65; variegated testa, 68-69
 Seedcoat, 130; correlation coefficients of yield, 105; and seed dormancy 65, 644; and herbicide injury, 241; and maturity, 626, 632; permeability of, 644; resistance of, 116, 506-507; splitting of, 115; starch accumulation, 415; and inner testa color, 69
 Seedcoat maggot, 523
 Seed/hull maturity index (SHMI), 628-630
 Seed isolates, 706-707
 Seedlings: abnormal vs. normal, 418; developmental needs, 415; and diseases of, 125, 362; growth of, 411 et passim.; and herbicide injuries, 241-242; morphology, 411; effect of refrigeration on vigor, 648; effect of season and location on vigor, 412
 Seedling feeder, 254F, 255
 Seedling rot, 375, 377
 Seed moisture content (SMC), 494-496
 Seed number: correlation coefficients, 75-76; general combining ability, 79; heterosis, 78; heritability estimates, 75-76; effect of temperature on, 441
 Seed parent, *A. hypogaea* as, 25, 29, 34-35
 Seed protectants, 364
 Seed rot, 364, 375, 377

- Seed splitter, 586, 588, 596T
 Seed stock, 604T-605T. *See also* Farmers stock peanuts
 Seed vigor tests, 418
 Segregating generations: correlations in, 75-77; genetic variability in, 75; heritability in, 75-77
 Segregation I, 576; and removal of aflatoxin-contaminated seed, 509; defined, 739; grading procedures, 508-509; seed types, 590; uses of, 592
 Segregation II, 576; defined, 739; grading procedures, 508; oilstock, 750-751; seed types, 590
 Segregation III, 576; and *A. flavus* infection, 499; defined, 739; grading procedure, 508-509; oilstock, 508, 750-751; seed types, 590
 Selection methods: accelerated programs, 102; bulk population or mass selection, 85, 100-102; genetic consequences of, 75-77; modified pedigree selection, 85, 102; pure line selection, 100, 102. *See also* Breeding
 Selection 65-121 (cultivar), 373
 Selfing, 28
 Semiochemicals, 762
 Sensible heat, 188, 190-191
 Sensitivity, 70-71
 Sensory quality: color, 670-671; flavor, 671-678; texture, 671; wholesomeness, 678-679
 Separation, 595; efficiency of, 592; rate of, 592
 Sequential branching, 50
 Serine, 656, 657T-659T, 700-701
 Serum cholesterol, 683
 Sesame extracts, 718
 Sesame flour, 726
 Sesone, 219-221, 241
 S-ethyl dipropylthiocarbamate (EPTC), 222-223
 S-(4,6-Diamino-s-triazin-2-ylmethyl)O-dimethylphosphorodithionate, 341
 Shading, 438-439
 Shaking, 458
 Shear thinning, 722
 Shelf life, 131
 Shell: importance of, 104-105; thin vs. thick, 64
 Shelled peanuts: and aflatoxin testing, 596, 598; sampling and grading, 595-596; standard grades, 597T; storage of, 600-602
 Sheller, 586, 592; effect of cylinder on seed quality, 413-414
 Shellers, peanut, 553; domestic market, 747-748; hedging techniques, 753; indemnification program, 509, 739; international market, 747, 749; lot assembling, 745; market risks, 751-752; processing and packaging, 745; and surplus peanuts for export, 738
 Shelling, 458, 624, 590-595; and curing treatments, 475; efficiency, 132, 592, 594-595; machines, 745; optimum design and settings, 610T-612T; output, 104; process, 608-613, 745; research needs, 762; sampling procedures, 508-509; seed damage and quality reduction, 413-414; and testa damage, 503-504
 Shelling outturn/percentage, 75, 595; combining ability, 79; and effects of drought, 173-174; inheritability, 73, 75; heterosis, 78; and insect damage, 530, 544; evaluation of stability, 84
 Shelling plants: insect infestation, 521-523, 541, 552; machines, 745; temperature extremes and insect control, 565
 Shellout Method, 626, 629, 632
 Shiny spider beetle, 522
 Shipping, 749
 Short position, 751-752
 Shrinkage, 574, 582
 Shrivelled pods. *See* "Raisins"
 Shulamit (cultivar), 110
 Siamese grain beetle, 521
 Sicklepod, 208T, 210F; date of germination, 207; response to herbicides, 222-223, 225, 232T; proclivity, 214, 215T, 216T, 218T, 244; effect on yields, 213-214
 Sida, 208T, 212F
 Sida, prickly: response to herbicides, 225, 232T; proclivity, 214, 215T, 216T, 218T
 Signalgrass, broadleaf, 208T; response to herbicides, 232T; proclivity, 214, 215T, 216T, 218T
 Silica gel-cellulose powder, 238
 Silt, percentage in soil, 238
 Silt loam soil, 228, 238-239
 Silty clay soil, 193
 Silvex, 220
 Sine cultivars, 62
 Single trailer drying systems, 482
 Skin effects, 229T
 Skinned seed outturns, 593
 Skin removal, 703
 Skin slippage, 475, 482-483
 Slag, 141
 Slenderhorned flour beetle, 521
 Slime disease, 376
 Sloane, Sir Hans, 4-5, 9-11, 14
 Slope-ratio technique, 679
 Smalleyed flour beetle, 521
 Smallflower morning glory. *See* Morning glory, smallflower
 Smartweed, Pennsylvania, 208T; response to herbicides, 232T; proclivity, 214

- Smith equation, 463, 465-466
 Smoke drying, 3-4
 Snack food, peanuts used for, 729-730, 764
 Soapstock, 510
 Soares de Souza, Gabriel, 3, 11, 14-15
 Sodium azide (SA), 70, 359
 Sodium chloride, 279, 510, 720
 Sodium content, 669T
 Sodium dodecyl sulfate (SDS), 696, 698
 Sodium hydroxide, 510
 Sodium hypochlorite, 511
 Sodium methylthiocarbamate, 356
 Sodium nitrate, 237
 Sodium salt, 224, 225, 227
 Sodium stearoyl-2-lactylate, 726
 Soil: acidity, 128, 141, 150, 237-238; analysis of, 242; fertility levels, 231, 237, 242, 412; effect on herbicides, 231-239; and irrigation practices, 193-195; and organic matter content, 238; effect of pH, 237-238, 242; optimum production requirements, 123-124; research needs, 756-757; temperature of, 236-237, 415; testing of, 128, 152; and water conductivity, 184; and water content, 184, 193, 195, 199. *See also* various types of soils
 Soil-borne diseases: aspergillus crown rot, 366-368; bacterial wilt, 376-377; black-hull, 371-372; botrytis blight, 374; charcoal rot, 368-369; cylindrocladium black rot, 357-359; diplodia collar rot, 369-371; fusarium diseases, 364-366; peanut clump, 345; pythium diseases, 359-362; research needs, 755; rhizoctonia diseases, 362-364; rhizopus diseases, 375; sclerotinia blight, 354-357; stem rot, 348-349, 352-354; verticillium wilt, 372-374; yellow mold, aflaroot, aflatoxin, 377-378
 Soil calcium. *See* Calcium
 Soil-inhabiting pests, 292-293; coleoptera, 301-305; and irrigation control, 201; lepidoptera, 293-301; resistance to, 132; white grubs, 305-307
 Soil moisture: and calcium uptake, 413; calibration of, 198; and digging time and losses, 459-460; and growth of diseases, 349, 361-362, 372, 376, 378; and fungi invasion, 492; effect on herbicides, 233-236; importance of, 242; monitoring of, 137; and stability of oil, 665; and transpiration rate, 192
 Soil surface moisture content: and evaporative demand, 190; and ground cover, 191; and pegging, 170-171
 Soil test B, 154
 Soil type: growth of diseases, 349, 367; and digging time, 459; effect on herbicides, 238-239; and mycoflora, 488, 490; and pest species and status, 258, 292-293; effect on seed quality, 412. *See also* various types of soils
 Soil water deficit (SWD): effect on evapotranspiration, 192; reduction in fruit addition, 183; partitioning to fruits, 169; reduction in flower numbers, 170; as an irrigation management strategy, 199; maintenance of leaf water content, 175; effect on root growth, 167; effect on vegetative growth, 168
 Soil water depletion (SWD): irrigation strategy, 195-197; and nitrogen fixation, 180; and vegetative growth, 168, 182
 Soil water measuring devices, 197-198
 Soil water potential, 174, 193; calibration of, 198; and irrigation, 195-197, 199; of different soil types, 194-195
 Soil water sensors, 196
 Solar energy drying systems, 483-484
 Solar radiation: and evapotranspiration, 187, 189; and disease control, 354; effect on growth rate, 438; as excess heat in storage, 582
 "Soldiers," 583
 Solubility, 707-712
 Soluble protein, 649
 Solvent classification, 702
 Solvent extraction, 510, 656, 673, 702
 Solvent recovery systems, 704
 Somatic chromosomes, 37-39
 Sorghum, 504. *See also* Grain sorghum
 Sorption processes, 463-466
 Sorting techniques, 595; for aflatoxin removal, 508-509; electronic, 618; rate of, 592
 Sound mature kernels, (SMK), 624, 497-501, 503; and AMI, 631; and bulk density, 579; defined, 588; and drought, 173-174
 Sound splits, 471, 483, 588
 Soup and beverage bases, peanuts used for, 723
 Southeastern Peanut Association, 595
 Southeastern production area, 258, 558
 Southeastern Runner 56-15; (cultivar), 280-281, 661
 Southern blight, 348, 755; and chemical control, 126, 329; and crop rotation, 126; and inadequate cultivation, 135; tolerance of, 43
 Southern corn rootworm: pest status, 258; and pod breakdown, 361; resistance to, 102, 111, 131-132; as a vector, 506
 Southern Regional Plant Introduction Station, 95
 Southwestern production area, 258, 558-559
 Soybean flours, 707

- Soybean meal, 486, 680
 Soybean oil, 682-683, 704, 723
 Soybean products, 702, 704, 717-718
 Soybeans, 175-177, 487, 755; bacterial control of insects, 566; comparison with peanut canopy, 207; in crop rotation, 125-126, 274, 389; and diseases, 339-340, 357, 359, 389; food value of, 750-751; and growth regulators, 135; response to herbicides, 223, 238-239, 241; and lipoxygenase activity, 637, 639; in meat products, 724; and pest species, 252; second order interactions, 83; vegetative and reproductive stages, 181; effects of water stress on AP, 423; weed competition in, 206-207; and yields, 147
 Soy isolates, 720
 Spacing, of peanut seed, 134. *See also* Row spacing
 Spangcross (cultivar): and AMI, 631; resistance of, 113, 388; yield increases, 104
 Spanhoma (cultivar), 115, 661-664
 Spanish erect (cultivar), 52, 73
 Spanish Improved (cultivar), 56, 60, 97
 Spanish jumbo (cultivar), 746
 Spanish type (cultivar), 14-15, 50, 53, 101, 419, 669T, 709; calcium requirement, 135; carbonyl compounds, 640, 674T; consumption patterns, 738; dormancy, 416, 642, 646; effects of drought, 169-171, 180, 182; early maturity breeding, 114; evapotranspiration, 185, 192; fatty acid composition, 115; genetic studies, 63, 74-78, 80, 86, 97, 100, 103; geographic areas of production, 123, 132-133, 738; and grade requirements, 586, 588, 595, 597F; effects of growth regulators, 435-438; irrigation scheduling, 197; effect of herbicides, 222, 228; kernel density, 628; response to liming, 141; market varieties of, 132-133; milling quality, 115; nitrogen and nitrogen fixation, 148-150, 180, 182; effect of photoperiod, 440; phenology and pest tolerance 254-256; and planting conditions, 134, 193; pod shape, 594F; resistance of, 102-103, 108-109, 112, 288, 334, 352, 361, 384; segregation of, 576; shelling and separating, 745; stability of oil, 114; susceptibility of, 367, 371, 373; volatile flavor components, 763
 Spanish-valencia cultivars, 181
 Spantex (cultivar), 74, 107, 132, 241
 Spatial tolerance, 255
 Specific combining ability (SCA), 79-81, 99-100
 Specific component quantitation, 625, 630-632
 Specific gravity, 595; removal of foreign material, 575; separation, 572, 606-608, 614, 616-618. *See also* Gravity tables
 Specific heat, 477-478
 Spectrophotometric maturity method, 627
 Spherosomes, 415
 Spider mites, 270, 280, 289-292; biology, 289; biological control of, 308T; damage, 290; management, 290-292; outbreaks of, 258-259; resistance to, 43; worldwide pest status, 309T
 Spider mite, two spotted, 330
 Spiders, 523
 Sphingolipid, 666
 Spindle inhibitor reagents, 37
 Spirea aphid, 344
 Split seed, 115; and fungi invasion, 488-492, 494-496, 499, 503-504; from insect damage, 530, 535, 544; and inverted windrow drying, 492; and shelling operation, 475; standards established for, 597T
 Split seed outturns, 593; and presizing, 613; and sheller speed, 608-613
 Sporulation, 496, 503
 Spotted wilt, 350F
 Spout sampler, 596T, 744
 Spray drying, 706, 709, 716-719, 723
 Spray insecticides, 279-280, 284, 288, 291, 298
 Spreader-sticker, 333
 Spreading growth habit, 52-53
 Sprinkler irrigation, 292
 S-propyl butylethylthiocarbamate, 222-223
 S-propyl dipropylthiocarbamate, 222-223
 Squarenecked grain beetle, 521
 Stability: of cultivars, 83-84; of oil, 84, 664-665
 Stabilization, 717-719
 Stachyose, 415, 667-668
 Stack cooker, 704
 Stackpole curing, 473-474; fungi invasion, 490-491, 498, 500; insect infestation, 552; effect on germinability, 413; loss of oil color, 636; damage to seed pod, 488
 Stackpole-stationary picker method, 116
 Standard crescent, 57-58
 Standard germination test, 418
 Starbur, bristly, 208T, 210F; response to herbicides, 232T; proclivity, 214, 215T, 216T, 218T
 Starch, 415, 644, 668

- Starr (cultivar), 74, 99, 132, 435F, 664; bulk density, 580T; ethylene-synthesis capacity, 643, 646; fatty acid composition, 661-662, 664; effect of herbicides and nematodes, 150, 241, 391; effect of irradiance and photoperiod, 439; milling quality, 592; response to nitrogen, 149; and nitrogen fixation, 150; and pod thickness, 594; resistance of, 107, 111, 113, 280-281, 288, 383, 388, 506; seedcoat splitting, 116; storage effects, 648; volatile compounds, 673, 674T-676T
 Starr spanish type, 425
 Static peanut pressures, 480, 579; and gravity tables, 617
 Stationary picker, 474
 Stearic acid, 660, 661T-663T, 682; mature vs. immature seed, 662; roasted vs. unroasted peanuts, 666T; triglycerides and phosphatides, 665T
 Steel alloys, 609
 Stem, peanut: and crescent factors, 58; inheritance of pigmentation, 63; inheritance of pubescence, 63; effect of soil water deficit on elongation, 168; optimum temperature for extension, 430
 Stem blight, 370
 Stem rot, 326, 348-354; causal organisms, 348-349, 362, 370; control of, 352-354; and dirtier cultivation, 219, 240; disease cycle and epidemiology, 349, 352; and nondirtier cultivation, 240; and ridge cultivation, 217; symptoms of, 348
 Sterigmatocystin, 512
 Sterile brachytic, 55-56
 Sterility, 58-59, 97-98
 Sterols, 666
 Sticks, as foreign material, 573T, 575T
 Stigmasterol, 666
 Sting nematodes, 379F, 383-384; and chemical control, 389-393; and crop rotation, 126; and lack of resistant cultivars, 388. *See also* Nematodes
 Stomata: activity of, 191; closure, 174-177, 193, 758; control, 183; frequency, 420; function, 178; effect of morphactins, 433; opening, 174-178, 192
 Storability, 575-576
 Storage, 574-586, 624, 635, 744-745; and artificial drying, 474; effects on dormancy, 646-647; effects on fatty acid composition, 665-666; and fungi invasion, 486, 492-496; effects on germination, 414, 647-650; and insect damage, 520, 523, 526, 528, 530, 539, 541; new methods of, 601-602; and milling process, 602; optimization of conditions in, 507-508; range of period, 576; degradation of protein, 416; and raw shelled peanuts, 600-602; research needs, 762; and volatiles, 640
 Storage atmospheres, 649. *See also* Moisture content
 Storage reserves, 415-416
 Storehouses. *See* Warehouse
 Strain evaluations, of rhizobium, 444-445. *See also* Rhizobia
 Strawberry sap beetle, 522
 Stress environments, 760
 Stubby root nematodes, 345, 384
 Stub-leaf, 412-413
 Stunt virus (PSV), 343; causal agent, 343-344; control of, 344; epidemiology, 344; resistance to, 43; symptoms and host range, 343
 Substituted anilines, 217
 Substomatal cavities, 190
 Subterranean pests. *See* Soil-inhabiting pests
 Succinic acid (2,2 dimethyl-hydrazide), 168, 433
 Succinic anhydride, 712-713, 716
 Succinylation, 710, 711F, 712, 716; and oil absorption, 721
 Sucrose, 171, 183, 415, 666-668, 722; and browning reaction, 671-672; effect of roasting, 669T
 Sugar amine reaction products, 671
 Sugarbeets, 207
 Sugar degradation products, 671
 Sugars, 146, 706; in after-ripening, 644; content, 415; effect of curing, 636
 Sulfides, 677T
 Sulfur: concentrations in whole plant, 154; critical levels of, 154; deficiency of, 153-154; in globulins, 655; sources, 153; uptake pattern, 154; and yield response, 153-154
 Sulfur amino acids, 43, 115, 679
 Sulfur compounds, 675T
 Sulfur dioxide, 510
 Sulfuric acid, 153, 157
 Sunflower seeds, 566; extractors, 718, flour, 726; oil, 660
 Supplemental irrigation, 137
 Suppression of insect pests, 252-253. *See also* Insect control, Pesticides
 Surface sprays, 557-558
 Surface treatments, 554
 Surge bins, 607
 Surge hoppers, 610-612T, 613
 Sweet clover, 333
 Sweet potatoes, 176; flour, 729
 Syneresis, 723

- Synergistic phenolic compounds, 643, 650
 Synergized pyrethrins, 554, 556, 562; resistance to, 564-565; schedule of treatment, 558-559
 Synergized pyrethrins supplemental surface sprays, 558-559
 Systems approach: to irrigation, 201; to weed control, 242-243
 Systems modeling, 257
 "Systems of weed control," 244
- ## T
- Tabares, Bishop of Valencia, 14
 Table salt: and aflatoxin reduction, 510; relative toxicity of, 230T
 Tall morning glory. *See* Morning glory, tall
 Tainan 6 (cultivar), 114
 Tainan 10 (cultivar), 114
 Tamnut (cultivar), 592, 594
 Tamnut 74 (cultivar), 132, 632
 Tanks, storage, 576
 Tannins, 68, 670; inhibition of *A. flavus*, 378, 507; leafspot resistance, 107; and increased permeability, 414; and PMI, 633
 Tarapota (cultivar), 108, 659T
 Taxonomic classification, 24-25, 42
 Taxonomic synthesis, 21-23
 T-bar commercial-type sheller, 609, 610T-611T
 TBR [V₄], 71
 TCMTB, 364
 Telone II, 389
 Temperature, 187, 206, 481-482; effect of altitude, 441; during blanching, 635; as a control measure, 354, 565; effects on drying, 467-469, 475-479; effect on evapotranspiration, 190; and functional properties of peanut protein, 708, 713; and growth of fungi, 487, 492, 494-497, 501-502, 504; effect on growth, 430-431, 441-443; effect on herbicide activity, 236-237; optimum for insect infestation, 526, 535, 543-544, 547, 549, 551-552; and irrigation scheduling 206; and determination of maturity, 625; effects on milling quality, 593-594; and equilibrium moisture content, 463, 475; effect on nitrogen fixation, 446; optimum for growth of pathogens, 328, 333, 338, 355, 358, 361, 367, 369-370, 372-375, 378; interaction with photoperiod, 440; interaction with relative humidity, 441; research needs, 760; and respiratory rates, 635; and specific heat values, 477-478; and stability of oil, 665; effects on storage, 574-575, 582, 584, 641, 646-648; effects on symbiosis, 446; ventilation and aeration systems and, 581
 Tempering room, 601
 Temporal tolerance, 254-255, 271
 Tenebrionids, 301
 Tennessee Red (cultivar), 116, 658T, 667
 Tensiometers, 137, 165, 197-198
 Tephrosia, 344
 Teratogenicity, 229
 Termites: and pod damage, 503, 506; worldwide pest status, 309T
 Terpenes, 675T
 Terretonin, 513
 Testa, 487; classes of, 66; color of, 65-68, 670; crescent factors and, 58; and crop maturity, 632; damage to, 503-504; integrity of, 116; and pod maturity profile, 634
 Tetrachloroethylene, 562
 Tetrachloroisophthalonitrile, 329
 Tetraethylthiuramdisulfide, 645
 Tetragenic inheritance, 55
 Tetrahydro-3,5-dimethyl-2H-1,3,5-thiadiazine-2-thione, 390
 Tetranychids, 289
 Tetrazolium, 618
 Texas panicum. *See* Panicum, Texas
 Texture, 671
 Textured vegetable protein, 724
 TG-1 (Vikram), 97
 Therevids, 297
 Thermal-type generators, 562-563
 Thermocouple psychrometry, 179
 Thermostat, 477
 Thiamine, 670, 684
 Thiamine hydrochloride, 447
 Thiamine monophosphate, 670
 Thiamine pyrophosphate, 670
 Thiazoles, 675T, 677T
 Thin-layer chromatographic technique (TLC), 598, 673
 Thin-layer drying relationships, 475
 Thiocarbamates, 217, 222-223
 Thiocyanogen value, 660T
 Thiol-reducine agents, 693, 698
 Thiopenes, 673, 677T
 Thiophanate, 372
 Thiram, 368-369, 645
 3-amino-2,5-dichlorobenzoic acid, 222
 3-methylbutanal, 673
 3-methyl-2-crotonolactone, 673
 3-(3,4-dichlorophenyl)-1-methoxy-1-methylurea, 222
 3-(3,4-dichlorophenyl)-1,1-dimethylurea, 220
 Threonine, 665, 657T-659T, 672, 679, 701
 Thrips, 270, 285; biology, 285-286; control of, 287-289, 308T, 391; damage, 286-287; genotype x pesticide interaction, 113; resistance to, 43, 106, 112; as a vector, 287; worldwide pest status, 309T

- TIBA, 433; effect on germination and vigor, 437; inhibition of peg elongation, 434; effect on peg strength, 435
 Tifrun (cultivar), 132, 144
 Tifspan (cultivar): rate of apparent photosynthesis, 426; evapotranspiration and, 185; soil water potential and irrigation scheduling, 196F, 197; effect of water stress, 174; yield response, 186F
 Tifton 8 (cultivar), 421
 Tifton loamy sand, 111, 113, 238-239
 Tillage, 242-243, 251. *See also* Cultivation
 Timing: of combining operations, 470-471; of digging operations, 458-460
 Tindivanam Variety-2, 642-643
 Tindivanam Variety-3, 642-643
 Tissue culture, 447; reproductive, 448-449; vegetative, 447-448
 Titer value, 660
 Tlalcacauatl, 4
 TMV 2, 199, 337, 642-643
 TMV 3, 642-643
 TMV 7, 337
 TMV 11, 337
 Toalson (cultivar), 109-110
 Tobacco, 79, 83, 125-126, 359
 Tobacco thrips, 43, 106, 131, 286
 Tocopherol composition, 665
 Tocopherols, 670, 683
 Tolerance, 254F; spatial, 255; temporal, 254-255
 Tomato cultivar Rutgers, 386
 Tomato spotted wilt, 341-342; causal agent, 342; control of, 342-343; epidemiology, 342; symptoms and host range, 342
 Tomato spotted wilt virus (TSWV), 282, 342; vectors of, 287-289
 Tomato wilt, 373
 'Topcrop', 340
 Top-dressing, of soluble calcium, 144
 Topography, and weed control, 242
 Total plant growth, 427-428
 Total radiation, 438
 Toxicity, 229T, 230T
 Toxicology, of herbicides registered for peanuts, 228-231
 Toxins, 290
 TP 393, 662T, 664T
 TP 716-2-1, 662T, 664T
 Tractor-powered combines, 552
 Tractors, and injury to plants, 355-356
 Transgressive segregation, 65
 Translocation: of calcium, 142; of herbicides, 226-228
 Transpiration, 174, 191-192; defined, 183; energy balance considerations and, 187; needs, 167; and soil water deficit, 192, and rapid changes in solar radiation, 189
 Transpiration, and milling process, 602
 Treflan, 225T
 Tremorgenic toxin, 512
 Triacylglycerol stereospecific structure, 633
 Tricarboxylic acid cycle, 415
 Trichloroacetic acid, 630
 Trichloronitromethane, 389
 Trichothecene mycotoxins, 513
 Triethanolamine salt, 226
 Trifluralin, 222, 224, 237-241; degradation of, 235; nomenclature, 225T; interaction with pesticides, 240; effect of lime, 238; relative toxicity of, 230T; residual activity, 236; soil moisture requirements, 233-234; effects of soil temperature, 236; technical description, 228; volatilization, 235; weeds controlled by, 232T
 Triglycerides, 660; and curing, 636, fatty acid composition, 665T
 Trilinolein, 639
 Trimethylpyrazine, 672
 Triphenyltin hydroxide, 329
 Tristimulus colorimetry, 671
 Troup loamy sand, 147
 Trypsin, 710, 716-717, 720
 Trypsin inhibitors, 704-705
 Tryptic proteolysis, 680
 Tryptophan, 43, 656, 657T, 681
 Tryptose, 722
 Tungsten carbide, 609
 Turbidity, 723
 Twobanded fungus beetle, 532
 21008, 288
 21016, 288
 21018, 288
 2-butanone, 636
 2-chloroethylphosphonic acid (CEPA), 417, 434, 645
 (2-chloroethyl) trimethylammonium chloride (CCC), 433, 436
 2-chloro-4-(ethylamino)-6-(isopropylamino)-S-triazine, 353
 2-crotonolactone, 673
 2,5-dimethylpyrazine, 672, 674T, 678T
 (2,4-bis(isopropylamino)-6-methylthio)-S-triazine, 222
 2,4-D, 219, 447-449, 645
 2,4-DB, 224, 241; nomenclature, 225T; postemergence application, 223; relative toxicity of, 230T; technical description, 228; weeds controlled by, 232T
 2,4-DEP, 219, 221
 2,4-dichlorophenoxyacetic acid (2,4-D), 219, 447-449, 645
 2,4-dinitrophenylhydrazones derivatives, 636
 2,4,5-T(4-amino-3,5,6-trichloropicolinic acid), 434
 (2,4,5-trichlorophenoxy)acetic acid (2,4,5-T), 219, 434

- 2-isopropyl-4,5-dimethylthiazole, 673
 2-methylbutanal, 673
 2-methylpropanal, 673
 2-methylpyrazine, 672
 2-methyl-2-(methylthio)propionaldehyde
 O-(methylcarbamoyl) oxime, 391
 2-nonanal, 636-637
 2-propyl-4,5 diethylthiazole, 673
 2-*sec*-butyl-4,6-dinitrophenol, 353
 2,6-dichloro-4-nitroaniline, 356
 2,6-dimethylpyrazine, 672, 674T, 678T
 2-*tert*-butyl-4-(2,4-dichloro-5-isopropoxyphenyl)- Δ^2 -1,3,4-oxadiazon-5-one, 353
 2-(thiocyanomethylthio)benzothiazole, 364
 2,3-dihydro-2,2-dimethyl-7-benzofuranyl methylcarbamate, 391
 2,3,5-trichloro-4-pyridinol, 222, 241
 2,3,5-triiodobenzoic acid, 433. *See also* TIBA
 (2-(2,4-dichlorophenoxy)ethyl)phosphite (2,4-DEP), 219, 221
 2-(2,4,5-trichlorophenoxy)propionic acid, 220
 Two spotted spider mite, 258, 330
 "Turkey X disease," 512
 Tyrosine, 655, 657T-659T, 700-701; and flavor, 672

U

- U-4-47-7 (EC 21115), 367
 Ultrafiltration, 706, 720-721
 Ultraviolet spectrophotometric method, 673
 United States: domestic food use, 655; as an exporter, 742, 750; first introduction of peanut, 14-15; germplasm collections, 95-96; harvesting and curing operations, 458; effect on international market, 749; and price support system, 571; production, in, 123, 124F, 737-740, 750-751, 764
 United States Department of Agriculture (USDA): administration of peanut program, 740; establishment of grading qualities, 508, 586, 739; standards for aflatoxin testing, 598T
 USDA Grade and State Seed Certification, regulations, 603
 U. S. Fancy, 599
 United States government: exportation, 738; overseas promotion, 740; price support system, 738-740, 749; role of, 737-740; self-regulation and, 739
 U. S. Jumbo, 559
 United States Standard Grades, 599

- Unloading equipment, 580T
 Unsaponifiable matter, 660T
 Unsaturated fatty acids, 682
 Unsaturation level, 636
 Urea: as a fertilizer, 149; in iron absorption, 156
 Urethane, 609
 Uric acid, 529
 Uses of peanuts, 13, 16-18, 510, 655; research needs, 763; roasters-salters, 746-747; segregation I, II, and III, 739, 750; split seed, 745

V

- Vacuum distillation, 673
 Vacuum drying, 483
 Vacuum storage, 649
 Valencia type (cultivar), 15, 50, 97, 101, 435; cleaned inshell peanuts, 598-599; dormancy, 416, 642, 646; fatty acid composition, 115, 661; genetic studies, 80, 86, 99-100; genotype x phenotype correlations, 75-77; geographic areas of production, 123, 133; grade requirements and, 586; heterosis, 78; market varieties, 133; response to nitrogen, 149; effect of photoperiod, 440; recessive leaf shape, 62; resistance of, 109, 112, 288, 334, 352, 506; segregation of, 576; stability of oil, 114; susceptibility of, 367, 371, 373, 384
 Valencia A. *See* New Mexico Valencia A
 Valencia spanish, 167
 Valine, 657T-659T, 679, 701
 Vanillic acid, 643
 Vapor phase, 469
 Vapor pressure, and windrow drying, 467
 Variance studies, 80-81
 Varied carpet beetle, 522
 Variegated testa, 67-69
 Variety selection: conditions of, 129; runner market type, 132; spanish market type, 132-133; valencia market type, 133; virginia market type, 130-132
 Vascular wilt: causes, 359-360, 365, 372; resistance to, 361
 Vegetative growth: analysis of, 428-429; and growth regulators, 135, 433-434; effect of heat stress, 760; and nitrogen accumulation, 151; and soil calcium requirement, 146, 756; effect of temperature, 443; effect of water deficit, 168-169, 182, 760
 Vegetative stages, 181
 Vegetative tissue, culture of, 447-448
 Velvetleaf, 207
 Ventilation fans, 564, 583, 745
 Ventilation system, 580T, 581, 583-584

- Verbascose, 415, 668
 Vernacular names, 9, 16. *See also individual names*
 Vernam, 225T
 Vernolate, 224, 238-239; control of annual weeds, 222-223; detoxification of, 237-238; interaction with herbicides, 240-241; and irrigation, 223; nomenclature, 225T; interaction with pesticides, 240; soil moisture requirement, 233-234; effect of soil pH, 237; effect of soil temperature, 236; subsurface application, 223; technical description, 228; relative toxicity of, 230T; weeds controlled by, 232T
 Verticillium wilt, 372; causal organism, 373; control of, 373-374; disease cycle and epidemiology, 373; resistance to, 71, 110; symptoms of, 372-373
 VGP 1, 359
 Viability, effects of temperature and relative humidity, 648
 Vibrational screening, 553
 Vigor, 418; effect of relative humidity, 414; storage effects, 414, 650; effect of sub-freezing temperatures, 413
 Vikram (cultivar), 97
 Vine clipping, 458, 460
 Violaxanthin, 61
 Virescent (chlorophyll-deficient mutant), 59-61
 Virginia production area, 258, 558-559
 Virginia type (cultivar), 15, 50, 53, 101, 419, 709; and AMI, 631; calcium requirement, 129, 172-173; cleaned inshell peanuts, 598-599; consumption patterns, 738; daily water requirements, 135; dormancy, 417, 646; effects of drought, 172-173; early maturity breeding, 114; fatty acid composition, 115; genetic studies, 74-78, 80, 82-83, 86, 99-100; geographic areas of production, 123, 130-131; grade requirements, 586, 588, 595, 597T; effect of growth regulators, 435-438; irrigation scheduling, 200; market varieties of, 130-132; milling quality, 115, 592; and nitrogen, 148-150; effect of photoperiod, 440; and planting conditions, 133-134; pod shape, 594F; postharvest chemical treatments, 645-647; preizing, 613; resistance of, 108, 112, 288, 334, 352, 373; segregation of, 576; shelling and separating, 745; specific heat values, 478; stability of oil, 114; susceptibility of, 358-359, 367, 371; yield variation, 104
 Virginia Beit Dagan (V4), 86
 Virginia Bunch cultivars, 97, 241; and A. *flavus* contamination, 489; and effective heat units, 625; evapotranspiration and,

- 185; fatty acid composition, 661; growth habit of, 52; resistance of, 111; and water uptake, 167; inheritance of yield, 74. *See also individual numbers*
 Virginia 56-R, 679
 Virginia Bunch 67: amino acid composition, 658T; monosaccharides, 667; physiological maturity methods, 632-633; resistance of, 288, 300, 373, 388
 Virginia 70, 669T
 Virginia 71-347, 356
 Virginia 72R, 291, 661, 662T, 664T
 Virginia 81 Bunch, 131
 Virginia Runner cultivars, 97; growth habit, 52; inheritance of yield, 73-74
 Viruses, as insect control mechanism, 566
 Virus transmission, 281-282
 Viscosity, 660T, 713
 Vitamin A, 670
 Vitamin B complex, 670
 Vitamin C, 670
 Vitamin D, 670
 Vitamin E, 670, 683-684
 Vitamin K, 670
 Volatile flavor components, 763
 Volatile profile analysis, 641
 Volatiles, 633, 636-640
 Volatilization, 235-236

W

- Wagon drying, 478-479
 Warehouse beetle, 522
 Warehouses: bulk protectant treatments, 555, 557; design features, 580T; rate of deterioration of peanuts, 520; equipment in, 573; insect pests in, 521-523, 526-527, 535, 541, 543, 549, 552-553; insecticidal aerosol space treatments, 561-564; storage effects, 647; structural designs of, 579; types of, 576-579
 Warehouse storage, 744-745
 Waspada (cultivar), 15
 Water, 25; daily water requirements, 136; flux density, 184; holding capacity of soil, 200; infiltration rates, 201; irrigation, 136-137, 201; leaching of, 705; management, 201, 443; and protein recovery, 694; rainfall, 125, 137; research needs, 755, 757-758; seasonal water requirements, 136; shortages of, 757, 761; potentials of soil types, 194; sorption of, 720; supplies of, 201; surface, 137; uptake of, 165-167, 414; and yield response, 186-187. *See also* Moisture content, Soil moisture
 Water account, bookkeeping method for, 200
 Water binding, 720-721

- Water content. *See* Moisture
- Water deficit: effect on calcium uptake, 172-173; on crop processes, 165-183; on flowering, 169-170; on leaf gas exchange, 174-178; and leaflet folding, 178; effect on nitrogen fixation, 180; on partitioning to fruits, 168-169, 182-183; on pegging and pod development, 170-171, 183; on seed quality and germinability, 173-174; on root growth, 165-167; on vegetative growth, 168, 182
- Watering schedule, programmed, 136
- Water retention curves, 194F
- Water stress, 174-178; response of apparent photosynthesis, 422-423; and flowering, 170-171; and leaflet folding, 178, 419; effect on nitrogen fixation, 446; research needs, 760
- Water use efficiency (WUE): defined, 184; research needs, 757-758; and row spacing, 193; and yield responses, 197
- Water vapor gradient, 178
- Water vapor loss, and diffusive resistance, 174
- Watery hypocotyl, 413
- Weather conditions, 761; and digging time, 459; and harvesting, 625-626; effect on yields, 442. *See also* Climatic conditions, Environment conditions
- Webbing damage, 524, 525F, 526F, 528-529
- Web blotch, 332, 350F; causal organism, 332-333; control of, 333-334; disease cycle and epidemiology, 333; resistance to, 43, 108, 334; symptoms of, 332
- Webb-Pomerene Corporation, 751
- Weed competition, 207-208, 213, effects of row spacing, 134; effect of time and duration, 213
- Weed control: of broadleaf weeds, 135; acceptance of chemicals, 212-213; and crop rotation, 125-126; through cultivation, 135, 242-243; interrelationship with diseases, 217-219; herbicides used in, 219-228; labeling requirements of herbicides, 231; "non-dirtting" vs. dirtting cultivation, 217-219, 240; in plant spacing, 134; principles of, 134-135; and postharvest problems, 412; research needs, 758; systems approach, 242-243
- Weed fruit, as foreign material, 573T, 575T
- Weeds, 208T; competitive ability, 213-215; and digging losses, 460; ecological population shifts of, 243; analyses of flora, 242; effect on harvesting, 214; major U. S. species, 214, 215T, 216T, 217, 218T; summary of, 243-245; effect on yields, 213
- Weighing lysimeters, 184
- Weight/weight relationships, 628-629; and crop maturity, 625, 628-630
- Wet-basis moisture content, 461-462
- "Wet" gypsum, 145
- Wet heat, 680
- Wettable powder (WP) spray, 554-558
- Wetting method, of restoring moisture to seed, 593
- Wheat, 78, 513, 755; control of CBR, 359; residual activity of herbicides, 236; trypsin inhibitors, 680
- Wheat straw-cellulose triacetate, 238
- Whey, 706, 722-723
- Whey protein gel matrix, 720
- White bread, peanuts used for, 726-727
- White clover, 344
- Whiteflies, 346-347
- White grubs, 270, 301, 305; biology, 305-306; damage, 306-307; management, 307; soil type preference, 293; worldwide pest status, 309T
- White mold, 126, 348
- White Spanish (cultivar), 153
- Wholesomeness, 678-679
- Whole wheat bread, peanuts used for, 726-727
- Wilco I (cultivar), 662T, 664T
- Wild cucumbers, as foreign material, 573T
- Wild species, 39; artificial hybridization techniques, 101; breeding strategies, 44; drought tolerance, 43; as gene pool, 42, 95-96, 98, 101; meiosis, 34; analysis of oil content, 43, 114; photosynthetic rate, 104, 421-422; protein content, 43; research in, 755; resistance traits, 43, 71, 292, 447; survival mechanisms, 416; tissue culture techniques, 450. *See also* Arachis
- Wind: and energy balance, 187; effect of evapotranspiration, 189, 191; movement of microorganisms, 487
- Windowpane fly, 523
- Windrow: and digging operations, 460; harvesting methods, 473-474; types of, 468; results of weathering, 595
- Windrow-combine method, 116
- Windrow drying/curing: aboveground field losses and, 470; and fungi invasion, 377, 490-492, 498-500, 513; effect on germinability, 413; and insect infestation, 552; rates of, 467, 488; research needs, 761
- Winged aphids, 341
- Wing petal shape, 58
- Winter plowing, 301
- Wireworms, 301
- "Worm-cut" seed damage, 524, 526F, 528, 529, 535, 544, 565
- WP-3, 681

X

- Xantha (radiation-induced mutant), 61
- Xanthamulata (X_m) (chemically induced mutant), 61
- Xanthophylls, 61
- X-ray induced sterile dwarfs, 56
- X-rays, in mutation production, 97
- Xylem sap expression, 179
- Xylem tissue, and calcium translocation, 142
- Xylem transpiration stream, 171, 172
- Xylene, 581
- Xylose, 668

Y

- Yields: effect of fertilization, 146-147, 154; genetic effects, 72 et passim. 80, 99-100; vs. growth period, 459F; and irrigation, 136, 165, 197; and planting date, 133;

- response to pesticides, 240, 393; effect of row spacing, 134; and soil calcium levels, 128-129, 139-141; in the U. S., 164; effect of weeds, 213-214, 219, 239
- Yield stability, 105-106, 755
- Yield variation, analysis of, 431-433
- Ynchic (peanut synonym), 1, 9, 16
- Yogurt systems, peanut used for, 722-723

Z

- (Z,E)-9,12-tetradecadien-1-ol acetate, 294
- Zinc: deficiencies of, 158; direct and residual effects, 158; foliar application, 158; importance of, 129; toxicity of, 158
- Zinc content, 668, 669T; of low quality seed, 413
- Zinc ion, and manganese ethylenebis-dithiocarbamate, 329
- Zinc sulfate, and suppression of sclerotinia blight, 356
- Zonate leafspot, 337